arXiv:1212.1815v1 [math.AG] 8 Dec 2012

ON THE COMPLEMENTS OF 3-DIMENSIONAL CONVEX POLYHEDRA AS POLYNOMIAL IMAGES OF $\mathbb{R}^3$

JOSÉ F. FERNANDO AND CARLOS UENO

Abstract. We prove that the complement $\mathcal{S} := \mathbb{R}^3 \setminus \mathcal{K}$ of a 3-dimensional convex polyhedron $\mathcal{K} \subset \mathbb{R}^3$ and its closure $\overline{\mathcal{S}}$ are polynomial images of $\mathbb{R}^3$. We also prove that for $n \geq 3$ there are bounded convex n-dimensional polyhedra $\mathcal{K} \subset \mathbb{R}^n$ such that none of its projections coincides with none of its sections, that is, for every affine projection $\pi : \mathbb{R}^n \to \mathbb{R}^n$ in the direction of a line $\vec{\ell}$ onto a hyperplane $H \subset \mathbb{R}^n$ the image $\pi(\mathcal{K})$ and the section $\mathcal{K} \cap H$ do not coincide. This property implies that the former techniques developed to represent complements $\mathcal{S}$ of unbounded convex polyhedra and their closures $\overline{\mathcal{S}}$ as regular images of $\mathbb{R}^n$ cannot be extended in general to represent such semialgebraic sets $\mathcal{S}$ and $\overline{\mathcal{S}}$ as polynomial images of $\mathbb{R}^n$ if $n \geq 4$.

Introduction

A map $f := (f_1, \ldots, f_m) : \mathbb{R}^n \to \mathbb{R}^m$ is a *polynomial map* if its components $f_k \in \mathbb{R}[\mathtt{x}] = \mathbb{R}[\mathtt{x}_1, \ldots, \mathtt{x}_n]$ are polynomials. Analogously, f is a *regular map* if its components can be represented as quotients $f_k = g_k/h_k$ of two polynomials $g_k, h_k \in \mathbb{R}[\mathtt{x}]$ such that h_k never vanishes on $\mathbb{R}^n$. The effective representation, if possible, of a subset $\mathcal{S} \subset \mathbb{R}^m$ as a polynomial or regular image of $\mathbb{R}^n$ reduces the study of certain classical problems in Real Geometry like optimization or Positivstellensätze results to its study in $\mathbb{R}^n$, with the clear profits that this involves (see [FG2, FU2]). Of course, polynomial representations are advantageous to regular representations, mainly in what concerns optimization, because they avoid denominators. During the last decade we have approached the problem of determining which (semialgebraic) subsets $\mathcal{S} \subset \mathbb{R}^m$ are polynomial or regular images of $\mathbb{R}^n$ (see [Fe, FG1, FG2, FGU, FU1, FU2, U1, U2, U3] for further details). By Tarski-Seidenberg's principle (see [BCR, 1.4]) the image of an either polynomial or regular map is a semialgebraic set. As it is well-known a subset $\mathcal{S} \subset \mathbb{R}^n$ is *semialgebraic* when it has a description by a finite boolean combination of polynomial equations and inequalities, which we will call a *semialgebraic* description.

A very distinguished family of semialgebraic sets is the one constituted by those semialgebraic sets whose boundary is piecewise linear, that is, semialgebraic sets which admit a semialgebraic description involving just polynomials of degree one. Of course, many of them cannot be polynomial or regular images of $\mathbb{R}^n$, as one easily realizes reviewing the known general conditions that must satisfy a semialgebraic set to be either a polynomial or regular image of $\mathbb{R}^n$ (see [FG1, FG2, FU1] for further details); however, it seems natural to wonder what happens with convex polyhedra, their interiors (as topological manifolds with boundary), their complements and the complements of their interiors. Since the

---

Date: 31/08/2012.

2010 *Mathematics Subject Classification.* Primary: 14P10, 52B11; Secondary: 14Q99, 90C26, 52B55.

Key words and phrases. Polynomial and regular maps and images, convex polyhedra, first and second trimming positions, optimization, Positivstellensätze, sectional projections.

First authors is supported by Spanish GR MTM2011-22435, while the second is a external collaborator of this project. This article is based on a part of the doctoral dissertation of the second author, written under the supervision of the first.

1-dimensional case is completely determined in [Fe] we assume in what follows that the dimension is ≥ 2.

We proved in [FGU] that all n-dimensional convex polyhedra and their interiors are regular images of $\mathbb{R}^n$. Since many convex polyhedra are bounded and the images of nonconstant polynomial maps are unbounded, the suitable general approach in that case was to consider regular maps; concerning the representation of unbounded polygons as polynomial images, see [U3]. On the other hand, in [FU2], we prove that the complement $\mathbb{R}^n \setminus \mathcal{K}$ of a convex polyhedron $\mathcal{K} \subset \mathbb{R}^n$ which does not disconnect $\mathbb{R}^n$ and the complement $\mathbb{R}^n \setminus \operatorname{Int} \mathcal{K}$ of its interior are regular images of $\mathbb{R}^n$. If $\mathcal{K}$ is moreover bounded or has dimension $d < n$, then $\mathbb{R}^n \setminus \mathcal{K}$ and $\mathbb{R}^n \setminus \operatorname{Int} \mathcal{K}$ are in fact polynomial images of $\mathbb{R}^n$.

Our purpose here is to show that the former techniques developed in [FU2] still work to represent as polynomials images of $\mathbb{R}^3$ the semialgebraic sets $\mathbb{R}^3 \setminus \mathcal{K}$ and $\mathbb{R}^3 \setminus \operatorname{Int} \mathcal{K}$ if $\mathcal{K} \subset \mathbb{R}^3$ is a 3-dimensional unbounded convex polyhedron. In this vein the main result of this work is the following:

Main Theorem. *Let $\mathcal{K} \subset \mathbb{R}^3$ be a 3-dimensional unbounded convex polyhedron which does not disconnect $\mathbb{R}^3$. Then $\mathbb{R}^3 \setminus \mathcal{K}$ and $\mathbb{R}^3 \setminus \operatorname{Int} \mathcal{K}$ are polynomial images of $\mathbb{R}^3$.*

Recall that a convex polyhedron $\mathcal{K} \subset \mathbb{R}^n$ disconnects $\mathbb{R}^n$ if and only if it is a *layer* (see [FU2, 1.2]), that is, it is affinely equivalent to $[-a, a] \times \mathbb{R}^{n-1}$ for some $a \in \mathbb{R}$, which reduces to a hyperplane if $a = 0$. These layers are particular cases of *degenerated* convex polyhedra which do not have vertices; we will refer to convex polyhedra which have at least one vertex as *nondegenerated*.

Proof of the Main Theorem. In view of the proofs of [FU2, Thm. 4.1 (ii) & 6.1 (ii)] and [FU2, Rmks 4.3 & 6.3], to prove the Main Theorem it is enough to guarantee that given a 3-dimensional convex polyhedron which has at least one vertex there exists a *facet* $\mathcal{F}$ (face of maximal dimension) such that $\mathcal{K}$ can be placed in both *extreme and strong first* and *second trimming positions* with respect to $\mathcal{F}$. Once this is done the proof of the Main Theorem runs very similar to the one in [FU2, Thm. 4.1 (ii) & 6.1 (ii)], and we leave to the reader the precise details. $\square$

Let us recall now such definitions, first introduced in [FU2], which we slightly modify here in order to clarify our discussion.

Trimming positions with respect to a facet. Consider in $\mathbb{R}^n$ the natural fibration induced by the projection $\pi_n : \mathbb{R}^n \to \mathbb{R}^n$, $(x_1, \ldots, x_n) \mapsto (x_1, \ldots, x_{n-1}, 0)$. The fiber $\pi_n^{-1}(a, 0)$ of π_n is a parallel line to the vector $\vec{e}_n := (0, \ldots, 0, 1)$ for each $a \in \mathbb{R}^{n-1}$. Given an n-dimensional convex polyhedron $\mathcal{K} \subset \mathbb{R}^n$, the intersection $\mathcal{I}_a := \pi_n^{-1}(a, 0) \cap \mathcal{K}$ can be either empty or a closed interval (either bounded or unbounded). Let $\mathcal{K} \subset \mathbb{R}^n$ be an n-dimensional convex polyhedron and let $\mathcal{F}$ be one of its facets. Consider the set $\mathfrak{A}_{\mathcal{K}} := \{a \in \mathbb{R}^{n-1} : \mathcal{I}_a \neq \varnothing,\ (a, 0) \notin \mathcal{I}_a\}$. We say that:

(1) $\mathcal{K}$ is in *first trimming position with respect to the facet* $\mathcal{F}$ if[1]:

- (i) $\mathcal{F} \subset \{x_{n-1} = 0\}$ and $\mathcal{K} \subset \{x_{n-1} \leq 0\}$.
- (ii) For all $a \in \mathbb{R}^{n-1}$ the interval $\mathcal{I}_a$ is bounded.
- (iii) The set $\mathfrak{A}_{\mathcal{K}}$ is bounded.

(2) $\mathcal{K}$ is in *second trimming position with respect to the facet* $\mathcal{F}$ if[2]:

---

[1] This definition corresponds to strong and extreme first trimming position in [FU2]

[2] This definition corresponds to strong and extreme second trimming position in [FU2]

(i) $\mathcal{F} \subset \{x_n = 0\}$ and $\mathcal{K} \subset \{x_n \leq 0\}$.

(ii) The set $\mathfrak{A}_{\mathcal{K}}$ is bounded.



FIGURE 1. (a) and (b) show convex polygons in second trimming position, and (c) shows a convex polyhedron in first trimming position.



FIGURE 2. (a) and (b) show convex polygons in second trimming position, and (c) shows a convex polyhedron in second trimming position.

Remarks. Let $\mathcal{K}\subset\mathbb{R}^n$ be an n-dimensional convex polyhedron and let $\mathcal{F}$ be one of its facets.

(i) If $\mathfrak{A}_{\mathcal{K}}$ is bounded, there is $N>0$ such that $A:=\bigcup_{a\in\mathfrak{A}_{\mathcal{K}}}\mathcal{I}_a\subset\mathfrak{A}_{\mathcal{K}}\times[-N,N]$, where $\mathcal{I}_a:=\pi_n^{-1}(a,0)\cap\mathcal{K}$ (see [FU2, 3.3, 5.2] for further details).

(ii) Suppose that $\mathcal{F}\subset\{x_n=0\}$ and $\mathcal{K}\subset\{x_n\leq 0\}$. For each $M\geq 0$ denote $\mathcal{K}_M:=\mathcal{K}\cup]-M,M[^n$. Then $\mathcal{K}$ is in second trimming position with respect to $\mathcal{F}$ if and only if there exists $M\geq 0$ such that $\pi_n(\mathcal{K}_M)=\mathcal{K}_M\cap\{x_n=0\}$.

The developement of a strategy to place (by means of an affine change of coordinates) an unbounded convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ in first or second trimming position with respect to a facet requires a good understanding of these positions in order to minimize the required operations to achieve it. Corollaries 2.4 and 2.5 go in the direction of providing a criterion to determine "easily" if a convex polyhedron is in first or second trimming position. Using them we get, for the 3-dimensional case, the following fully satisfactory placing results, which provide the proof of the Main Theorem. Namely,

Theorem 1 (3-dimensional polyhedra and first trimming position)**.** *Let $\mathcal{K}\subset\mathbb{R}^3$ be a 3-dimensional unbounded convex polyhedron. Then*

(i) *The convex polyhedron $\mathcal{K}$ can be placed in first trimming position with respect to some of its unbounded facets $\mathcal{F}$.*

(ii) *Moreover, if all the unbounded edges of $\mathcal{K}$ are parallel, then $\mathcal{K}$ has at least one bounded facet and it can be placed in first trimming position with respect to any of its bounded facets.*

Remark. From the proof of Theorem 1 we can deduce that, in fact, $\mathcal{K}$ can be placed in first trimming position with respect to any facet with nonparallel unbounded edges.

Theorem 2 (3-dimensional polyhedra and second trimming position)**.** *Let $\mathcal{K}\subset\mathbb{R}^3$ be a 3-dimensional unbounded convex polyhedron. We have:*

(i) *If $\mathcal{K}$ has facets with nonparallel unbounded edges, then it can be placed in second trimming position with respect to any of them.*

(ii) *If all the unbounded edges of $\mathcal{K}$ are parallel, then $\mathcal{K}$ can be placed in second trimming position with respect to any of its bounded facets.*

The obsructions to extend the Main Theorem to dimension ≥ 4, using our techniques, concentrates on the fact that the following property is exclusive of convex polygons.

Property. *Let $\mathcal{K}\subset\mathbb{R}^n$ be an n-dimensional convex polyhedron and let $\vec{\ell}\subset\mathbb{R}^n$ be a vectorial line; consider the canonical projection $\pi:\mathbb{R}^n\to\mathbb{R}^n/\vec{\ell}$. Then there is a hyperplane H of $\mathbb{R}^n$ such that $\pi(\mathcal{K})=\pi(\mathcal{K}\cap H)$.*

In Theorem 4.2 we prove that for $n\geq 3$ there are n-dimensional bounded convex polyhedra $\mathcal{K}\subset\mathbb{R}^n$ *without sectional projections*, that is, for each line $\vec{\ell}$ and each hyperplane H the projection $\pi:\mathbb{R}^n\to\mathbb{R}^n$ in the direction of $\vec{\ell}$ onto the hyperplane H satisfies $\pi(\mathcal{K})\neq\mathcal{K}\cap H$. This allows us to produce examples of n-dimensional unbounded convex polyhedra of the type $\mathcal{K}\times[0,+\infty)$ (with $\mathcal{K}$ as above) that can be placed neither in first trimming position with respect to any of its facets nor in second trimming position with respect to any unbounded facet.

The article is organized as follows. In Section 1 we present the main concepts and tools related to convex polyhedra that will be used in the subsequent sections. In Section 2 we analyze the sectional projections of a polyhedron and we provide the tools to approach the proofs of Theorems 1 and 2, which are developed in Section 3. Finally in Section 4, we prove Theorem 4.2 concerning the existence of convex polyhedra with no sectional projections.

1. Convex polyhedra

We begin by introducing some preliminary terminology and notations concerning convex polyhedra. For a detailed study of the main properties of convex sets, we refer the reader to [Be1, Be2, R]. Along this work, we write $\mathbb{R}[\mathtt{x}]$ to denote the ring of polynomials $\mathbb{R}[\mathtt{x}_1,\dots,\mathtt{x}_n]$. Given an affine hyperplane $H\subset\mathbb{R}^n$ there exists a degree one polynomial $h\in\mathbb{R}[\mathtt{x}]$ such that $H:=\{x\in\mathbb{R}^n:\ h(x)=0\}\equiv\{h=0\}$. This polynomial h determines two *closed half-spaces* H^+ and H^-, defined by

$$H^+:=\{x\in\mathbb{R}^n:\ h(x)\geq 0\}\equiv\{h\geq 0\}\quad\text{and}\quad H^-:=\{x\in\mathbb{R}^n:\ h(x)\leq 0\}\equiv\{h\leq 0\}.$$

In this way, the labels H^+ and H^- are assigned in terms of the choice of h (or equivalently, in terms of the orientation of H) and it is enough to substitute h by $-h$ to interchange the roles of H^+ and H^-. As usual, we denote by $\vec{h}$ the homogeneous component of h of degree one and by $\vec{H}:=\{\vec{v}\in\mathbb{R}^n:\ \vec{h}(\vec{v})=0\}$ the *direction of* H. More generally, we also denote by $\vec{W}$ the direction of any affine subspace $W\subset\mathbb{R}^n$. As usual, we write $\mathcal{B}_n(p,r)$ for the open ball of center $p\in\mathbb{R}^n$ and radius $r>0$, and $\overline{\mathcal{B}}_n(p,r)$ for the corresponding closed ball. The boundary of such a ball is the $(n-1)$-dimensional sphere $\mathbb{S}^{n-1}(p,r)$ of center $p\in\mathbb{R}^n$ and radius $r>0$.

1.1. Generalities on convex polyhedra.

A subset $\mathcal{K}\subset\mathbb{R}^n$ is a *convex polyhedron* if it can be described as the finite intersection $\mathcal{K}:=\bigcap_{i=1}^r H_i^+$ of closed half-spaces H_i^+; we allow this family of half-spaces to by empty to describe $\mathcal{K}=\mathbb{R}^n$ as a convex polyhedron. The dimension $\dim(\mathcal{K})$ of $\mathcal{K}$ is its dimension as a topological manifold with boundary, which coincides with its dimension as a semialgebraic subset of $\mathbb{R}^n$ and with the dimension of the smallest affine subspace of $\mathbb{R}^n$ which contains $\mathcal{K}$.

1.1.1. Let $\mathcal{K}\subset\mathbb{R}^n$ be an n-dimensional convex polyhedron. By [Be2, 12.1.5] there exists a unique minimal family $\mathfrak{H}:=\{H_1,\dots,H_m\}$ of affine hyperplanes of $\mathbb{R}^n$, which is empty just in case $\mathcal{K}=\mathbb{R}^n$, such that $\mathcal{K}=\bigcap_{i=1}^m H_i^+$; we refer to this family as the *minimal presentation* of $\mathcal{K}$. Of course, we will always assume that we have chosen the linear equation h_i of each H_i such that $\mathcal{K}\subset H_i^+$.

The *facets* or $(n-1)$-*faces* of $\mathcal{K}$ are the intersections $\mathcal{F}_i:=H_i\cap\mathcal{K}$ for $1\leq i\leq m$; only the convex polyhedron $\mathbb{R}^n$ has no facets. Each facet $\mathcal{F}_i:=H_i^-\cap\bigcap_{j=1}^m H_j^+$ is a polyhedron contained in H_i. For inductive processes, we will denote $\mathcal{K}_{i,\times}:=\bigcap_{j\neq i}H_j^+$, which is a convex polyhedron that strictly contains $\mathcal{K}$ and satisfies $\mathcal{K}=\mathcal{K}_{i,\times}\cap H_i^+$.

A convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ is a topological manifold, whose interior $\operatorname{Int}\mathcal{K}$ is a topological manifold without boundary, which coincides with the topological interior of $\mathcal{K}$ in $\mathbb{R}^n$ if and only if $\dim(\mathcal{K})=n$. The *boundary* of $\mathcal{K}$ is $\partial\mathcal{K}=\mathcal{K}\setminus\operatorname{Int}\mathcal{K}$; in particular, if $\mathcal{K}$ is a singleton, then $\operatorname{Int}\mathcal{K}=\mathcal{K}$ and $\partial\mathcal{K}=\varnothing$. By [Be2, 12.1.5], $\partial\mathcal{K}=\bigcup_{i=1}^m\mathcal{F}_i$, and so

$$\operatorname{Int}\mathcal{K}=\mathcal{K}\setminus\bigcup_{i=1}^m(\mathcal{K}\cap H_i)=\bigcap_{j=1}^m H_j^+\cap\bigcap_{i=1}^m(\mathbb{R}^n\setminus H_i)=\bigcap_{i=1}^m(H_i^+\setminus H_i).$$

For $0\leq j\leq n-2$, we define inductively the j-*faces* of $\mathcal{K}$ as the facets of the $(j+1)$-faces of $\mathcal{K}$, which are once more convex polyhedra. As usual, the 0-faces are the *vertices* of $\mathcal{K}$ and the 1-faces are the *edges* of $\mathcal{K}$; obviously if $\mathcal{K}$ has a vertex, then $m\geq n$ (see [Be2, 12.1.8-9]).

A convex polyhedron of $\mathbb{R}^n$ is *nondegenerated* if it has at least one vertex; otherwise, we say that the convex polyhedron is *degenerate.* In general, we denote by $\mathcal{E}$ a generic face of $\mathcal{K}$ to distinguish it from the facets $\mathcal{F}_1,\dots,\mathcal{F}_m$ of $\mathcal{K}$; the affine subspace generated by $\mathcal{E}$ is denoted by W to be distinguished from the affine hyperplanes $H_1,\dots,H_m$ generated by the facets $\mathcal{F}_1,\dots,\mathcal{F}_m$ of $\mathcal{K}$.

1.1.2. Observe that if $H\subset\mathbb{R}^n$ is a hyperplane we have $\partial H^+=H=\partial H^-$. A *supporting hyperplane* of a convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ is a hyperplane H of $\mathbb{R}^n$ which intersects $\mathcal{K}$ and satisfies $\mathcal{K}\subset H^+$ or $\mathcal{K}\subset H^-$. As one can check, the intersection of $\mathcal{K}$ with a supporting hyperplane H is a face of $\mathcal{K}$ and conversely, each face of $\mathcal{K}$ is the intersection of $\mathcal{K}$ with some supporting hyperplane. In particular, the vertices of a convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ are those points $p\in\mathcal{K}$ for which there is a (supporting) hyperplane $H\subset\mathbb{R}^n$ such that $\mathcal{K}\cap H=\{p\}$. Moreover, the supporting hyperplanes of $\mathcal{K}$ coincide with the hyperplanes $H\subset\mathbb{R}^n$ such that $\varnothing\neq\mathcal{K}\cap H\subset\partial\mathcal{K}$. All these facts will be used freely along this work.

1.1.3. An important property of a bounded convex polyhedron is that it coincides with the convex hull of the set of its vertices, that is, if $p_1,\dots,p_r$ are the vertices of $\mathcal{K}$, then

$$\mathcal{K}=\Big\{\sum_{i=1}^r\mu_ip_i:\ \mu_i\geq 0,\ \sum_{i=1}^r\mu_i=1\Big\}.$$

For a general convex polyhedron, there is a similar characterization which involves also the directions of the unbounded edges (see Corollary 1.8).

A particular type of convex polyhedra are the cones (bounded or unbounded). The *bounded cone of vertex p and base the bounded convex polyhedron $\mathcal{P}\subset\mathbb{R}^n$* is the set

$$\mathcal{C}:=\{\lambda p+(1-\lambda)q:\ q\in\mathcal{P},\ 0\leq\lambda\leq 1\}.$$

On the other hand, given vectors $\vec{v}_1,\dots,\vec{v}_r\in\mathbb{R}^n$ we define the *(vectorial) cone generated by the vectors $\{\vec{v}_1,\dots,\vec{v}_r\}$* as the set $\vec{\mathcal{C}}:=\{\sum_{i=1}^r\lambda_i\vec{v}_i:\ \lambda_i\geq 0\}$; now, the (unbounded) cone of vertex p generated by the vectors $\{\vec{v}_1,\dots,\vec{v}_r\}$ is $\mathcal{C}:=p+\vec{\mathcal{C}}$.

In what follows, given two points $p,q\in\mathbb{R}^n$ we denote by $\overline{pq}:=\{\lambda p+(1-\lambda)q:\ 0\leq\lambda\leq 1\}$ the *(closed) segment connecting p and q.* On the other hand, given a point $p\in\mathbb{R}^n$ and a vector $\vec{v}\in\mathbb{R}^n$, we denote by $p\vec{v}:=\{p+\lambda\vec{v}:\lambda\geq 0\}$ the *(closed) half-line of extreme p and direction $\vec{v}$.* Clearly, the boundary of the segment $\mathcal{S}:=\overline{pq}$ is $\partial\mathcal{S}=\{p,q\}$ and its interior is $\operatorname{Int}\mathcal{S}=\mathcal{S}\setminus\{p,q\}$, while the boundary of the half-line $\mathcal{H}:=p\vec{v}$ is $\partial\mathcal{H}=\{p\}$ and its interior is $\operatorname{Int}\mathcal{H}=\mathcal{H}\setminus\{p\}$.

1.1.4. The *affine maps* are those maps that preserve affine combinations; hence, affine maps are polynomial maps that preserve convexity and so they transform convex polyhedra onto convex polyhedra. As usual, the linear map (or derivative) associated to an affine map $f:\mathbb{R}^n\to\mathbb{R}^m$ is denoted by $\vec{f}:\mathbb{R}^n\to\mathbb{R}^m$. It will be common to use affine bijections $f:\mathbb{R}^n\to\mathbb{R}^n$, that will be called *changes of coordinates*, to place a convex polyhedron, or more generally a semialgebraic set $\mathcal{S}$, in advantageous positions. As usual we say that $\mathcal{S}$ and $f(\mathcal{S})$ are *affinely equivalent.*

1.2. Recession cone of a convex polyhedron.

We can associate to each convex polyhedron its *recession cone,* that we define next (see also [Z, Ch.1]).

Definition and Proposition 1.1. Let $\mathcal{K}\subset\mathbb{R}^n$ be a convex polyhedron. Fix a point $p\in\mathcal{K}$ and consider the set $\vec{\mathfrak{C}}(\mathcal{K}):=\{\vec{v}\in\mathbb{R}^n:\ p\vec{v}\subset\mathcal{K}\}$. *Then $\vec{\mathfrak{C}}(\mathcal{K})$ is a (vectorial)* *convex cone and it does not depend on the choice of* p. The set $\vec{\mathfrak{C}}(\mathcal{K})$ will be called the *recession cone* of $\mathcal{K}$.

Proof. To prove that $\vec{\mathfrak{C}}(\mathcal{K})$ is a convex cone it is enough to check that if $\vec{v},\vec{w}\in\vec{\mathfrak{C}}(\mathcal{K})$ and $\lambda,\mu\geq 0$, then $\lambda\vec{v}+\mu\vec{w}\in\vec{\mathfrak{C}}(\mathcal{K})$; the concrete details are left to the reader. To prove that $\vec{\mathfrak{C}}(\mathcal{K})$ does not depend on the point p, we verify that the sets

$$\vec{\mathfrak{C}}_p(\mathcal{K}):=\{\vec{v}\in\mathbb{R}^n:\, p\vec{v}\subset\mathcal{K}\}\quad\text{and}\quad\vec{\mathfrak{C}}_q(\mathcal{K}):=\{\vec{v}\in\mathbb{R}^n:\, q\vec{v}\subset\mathcal{K}\}$$

coincide. In fact, it is enough to check that given $\vec{v}\in\vec{\mathfrak{C}}_p(\mathcal{K})$, it holds that $q\vec{v}\subset\mathcal{K}$. We may assume that $q\notin p\vec{v}\cup p(-\vec{v})$; otherwise, using the convexity of $\mathcal{K}$ the inclusion $q\vec{v}\subset\mathcal{K}$ is clear.

Indeed, let $x\in q\vec{v}\setminus\{q\}$ and $\mu>0$ be such that $x-q=\mu\vec{v}$. Choose a sequence of points $\{y_k\}_{k\in\mathbb{N}}\subset p\vec{v}\setminus\{p\}$ such that the sequence $\{\|y_k-p\|\}_{n\in\mathbb{N}}$ tends to $+\infty$. For each $k\in\mathbb{N}$, let $\rho_k>0$ be such that $y_k-p=\rho_k\vec{v}=\lambda_k(x-q)$, where $\lambda_k:=\rho_k/\mu>0$. Note that also the sequence $\{\lambda_k=\|y_k-p\|/\|x-q\|\}_{k\in\mathbb{N}}$ tends to $+\infty$. Since $q,y_k\in\mathcal{K}$ and this polyhedron is convex, each segment $\overline{qy_k}$ is contained in $\mathcal{K}$. Now, since $y_k=p+\lambda_k(x-q)$, we have

$$z_k:=\Big(\frac{1}{1+\lambda_k}\Big)p+\Big(\frac{\lambda_k}{1+\lambda_k}\Big)x=\Big(\frac{\lambda_k}{1+\lambda_k}\Big)q+\Big(\frac{1}{1+\lambda_k}\Big)y_k\in\overline{qy_k}\subset\mathcal{K}.$$

Thus, since $\mathcal{K}$ is a closed subset of $\mathbb{R}^n$, we conclude

$$x=\lim_{k\to\infty}\Big(\Big(\frac{\lambda_k}{1+\lambda_k}\Big)x+\Big(\frac{1}{1+\lambda_k}\Big)p\Big)=\lim_{k\to\infty}z_k\in\mathcal{K},$$

as wanted. □

Remarks 1.2. Let $\mathcal{K}\subset\mathbb{R}^n$ be a convex polyhedron and let $p\in\mathcal{K}$.

(i) $\vec{\mathfrak{C}}(\mathcal{K})=\{\mathbf{0}\}$ if and only if $\mathcal{K}$ is bounded.

(ii) The set $\mathfrak{C}_p(\mathcal{K})=p+\vec{\mathfrak{C}}_p(\mathcal{K})$ is either equal to $\{p\}$ if $\mathcal{K}$ is bounded or the union of all the closed half-lines of $\mathbb{R}^n$ contained in $\mathcal{K}$ and whose boundary is $\{p\}$ if $\mathcal{K}$ is unbounded.

(iii) A straightforward computation shows that if $\mathcal{P}\subset\mathbb{R}^n$ is a nondegenerated convex polyhedron and $k\geq 1$, then $\vec{\mathfrak{C}}(\mathbb{R}^k\times\mathcal{P})=\mathbb{R}^k\times\vec{\mathfrak{C}}(\mathcal{P})$. Since degenerated convex polyhedra can be written, after a change of coordinates, as the product of a nondegenerated convex polyhedron times an Euclidean space (see [FGU, §2.2]), we only need to study the recession cones of nondegenerate convex polyhedra.

(iv) Given two convex polyhedra $\mathcal{K}_1\subset\mathcal{K}_2$, it holds $\vec{\mathfrak{C}}(\mathcal{K}_1)\subset\vec{\mathfrak{C}}(\mathcal{K}_2)$.

(v) Moreover, if $\mathcal{K}:=\bigcap_{i=1}^r H_i^+$, then $\vec{\mathfrak{C}}(\mathcal{K}):=\bigcap_{i=1}^r\vec{\mathfrak{C}}(H_i^+)=\bigcap_{i=1}^r\vec{H_i}^+$. Consequently, if $\{\mathcal{K}_1,\ldots,\mathcal{K}_s\}$ are convex polyhedra with nonempty intersection $\mathcal{K}:=\bigcap_{j=1}^s\mathcal{K}_j$, then $\vec{\mathfrak{C}}(\mathcal{K}):=\bigcap_{j=1}^s\vec{\mathfrak{C}}(\mathcal{K}_j)$.

1.3. Facing upwards positions for convex polyhedra.

As we have commented in the introduction, the proof of the Main Theorem reduces to show that nondegenerated convex polyhedra of $\mathbb{R}^3$ can be placed in a concrete form. As an initial step to achieve our aim we introduce the concept of *facing upwards position.*

Definitions 1.3. We say that an unbounded nondegenerated convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ is in *facing upwards position (or* FU*-position) with respect to the (vectorial and oriented) hyperplane* $\vec{\Pi}$ of $\mathbb{R}^n$ if there is a hyperplane Π parallel to $\vec{\Pi}$ which intersects all the unbounded edges of $\mathcal{K}$ and such that all the vertices of $\mathcal{K}$ belong to the open half-space

$\operatorname{Int}\Pi^-$. Such a hyperplane Π will be called a *sawing hyperplane* for $\mathcal{K}$. Of course, any other hyperplane $\Pi' \subset \Pi^+$ (necessarily parallel to Π) is also a sawing hyperplane for $\mathcal{K}$.

Now, if $\vec{h} \in \mathbb{R}[\mathtt{x}]$ is a (homogeneous) degree one polynomial describing $\vec{\Pi}$ and $h := 0+\vec{h}$, we say that the FU-position of $\mathcal{K}$ with respect to $\vec{\Pi}$ is *optimal* if the minimum of $h|_{\mathcal{K}}$ is attained just in one point (which must be a vertex of $\mathcal{K}$); such point is the *minimal vertex* of $\mathcal{K}$ (with respect to $\vec{\Pi}$).

Next, we prove that each nondegenerated convex polyhedron can be placed in optimal FU-position with respect to the hyperplane $\vec{\Pi} := \{x_n = 0\}$.

Lemma 1.4. *Let $\mathcal{K} \subset \mathbb{R}^n$ be an n-dimensional nondegenerated convex polyhedron. Then $\mathcal{K}$ can be placed in optimal* FU*-position with respect to the hyperplane $\vec{\Pi} := \{x_n = 0\}$ in such a way that $\mathcal{K} \subset \{x_n \geq -1\}$.*

Proof. Let $\{H_1, \dots, H_m\}$ be the minimal presentation of $\mathcal{K}$. Since $\mathcal{K}$ is nondegenerated, $m \geq n$ and we may assume after a change of coordinates that $\bigcap_{i=1}^n H_i = \{\mathbf{0}\}$ is a vertex of $\mathcal{K}$ and $H_i^+ = \{x_i \geq 0\}$ for $1 \leq i \leq n$. Consequently,

$$\mathcal{K} \subset \{x_1 \geq 0, \dots, x_n \geq 0\} \subset \{x_1 + \cdots + x_n > 0\} \cup \{\mathbf{0}\}.$$

Next, we apply to $\mathcal{K}$ a change of coordinates that transforms the half-space $\{x_1+\cdots+x_n \geq 0\}$ onto $\{h := x_n \geq 0\}$ and keeps fixed the vertex $\mathbf{0}$. Notice that $0 = h(\mathbf{0}) < h(x)$ for $x \in \mathcal{K} \setminus \{\mathbf{0}\}$. Let us see now that $\mathcal{K}$ can be placed in optimal FU-position with respect to the hyperplane $\vec{\Pi} := \{\vec{h} := x_n = 0\}$; obviosuly, $\mathcal{K} \subset \{x_n \geq -1\}$.

First, we choose a hyperplane $\Pi := \{x_n = M\}$ for some $M > 0$ such that all the vertices of $\mathcal{K}$ are contained in $\operatorname{Int}\Pi^-$. Let now $\mathcal{A}$ be an unbounded edge of $\mathcal{K}$. Since $\mathcal{K}$ is nondegenerated, it cannot contain lines (see [FG1, §2.2]) and so there exist a vertex $p := (p_1, \dots, p_n) \in \mathcal{K}$ and a vector $\vec{v} := (v_1, \dots, v_n) \in \vec{\mathfrak{C}}(\mathcal{K})$ such that $\mathcal{A} = p\vec{v}$; observe that $p_n < M$. Since $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})$, the half-line $\mathbf{0}\vec{v} \subset \mathcal{K}$ and since $\mathcal{K} \setminus \{\mathbf{0}\} \subset \{x_n > 0\}$, we have $v_n > 0$; hence, there exists $t_0 > 0$ such that $p_n + t_0 v_n = M$, that is, $p + t_0\vec{v} \in \mathcal{A} \cap \Pi$, and we are done. □

We will see in Remark 2.2 that the intersection of a nondegenerated unbounded convex polyhedron in FU-position with respect to a hyperplane $\vec{\Pi}$ and a sawing hyperplane Π for $\mathcal{K}$ is a bounded polyhedron $\mathcal{P} := \mathcal{K} \cap \Pi$ whose vertices are the intersections with Π of the unbounded edges of $\mathcal{K}$. Keeping this in mind we now show a way to construct a supporting hyperplane of $\mathcal{K}$ so that its intersection with $\mathcal{K}$ is an unbounded edge.

Lemma 1.5. *Let $\mathcal{K} \subset \mathbb{R}^n$ be a nondegenerated unbounded convex polyhedron in FU-position with respect to a hyperplane $\vec{\Pi}$ and let Π be a sawing hyperplane for $\mathcal{K}$. Let W be a supporting hyperplane of the polyhedron $\mathcal{P} := \mathcal{K} \cap \Pi$ in Π, let $p \in W \cap \mathcal{P}$ be a vertex of $\mathcal{P}$ and let $\mathcal{A}$ be the unbounded edge of $\mathcal{K}$ such that $\{p\} = \mathcal{A} \cap \Pi$. Then the affine subspace H of $\mathbb{R}^n$ generated by W and $\mathcal{A}$ is a supporting hyperplane of $\mathcal{K}$.*

Proof. First, H is a hyperplane of $\mathbb{R}^n$ with $H \cap \Pi = W$, because $\mathcal{A} \cap W = \{p\}$, $\dim \mathcal{A} = 1$ and $\dim W = n-2$. On the other hand, since $\mathcal{A} \subset H \cap \mathcal{K}$, to prove that H is a supporting hyperplane of $\mathcal{K}$, it is enough to prove that $H \cap \mathcal{K} \subset \partial\mathcal{K}$. We write $\mathcal{A} := p_0\vec{u}$, where p_0 is a vertex of $\mathcal{K}$ and $\vec{u} \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{\mathbf{0}\}$. Observe that $\mathcal{A}$ intersects $\operatorname{Int}\Pi^+$ because $p_0, p \in \mathcal{A}$, $p_0 \in \operatorname{Int}\Pi^-$ and $p \in \Pi$. We will use freely along the rest of the proof the following well-known property of convex sets (see [Be1, 11.2.4]):

(1.5.1) *Let* $S \subset \mathbb{R}^n$ *be a convex set with nonempty interior and let* $p \in \operatorname{Int}(S)$ *and* $q \in \operatorname{Cl}(S)$. *Then* $\overline{pq} \setminus \{q\} \subset \operatorname{Int}(S)$.

Suppose by way of contradiction that there is $q \in H \cap \operatorname{Int} \mathcal{K}$. Let us check first that $q \in \operatorname{Int} \Pi^+$. Otherwise, $q \in \Pi^-$; choose a point $y \in \mathcal{A} \cap \operatorname{Int} \Pi^+$ and observe that the segment $\overline{qy} \subset H$ meets Π at a point $b \in H$ different from y. Since $q \in \operatorname{Int} \mathcal{K}$, we have $b \in \overline{qy} \setminus \{y\} \subset \operatorname{Int} \mathcal{K}$ and so $b \in \Pi \cap \operatorname{Int} \mathcal{K} = \operatorname{Int} \mathcal{P}$; hence, $b \in H \cap \operatorname{Int} \mathcal{P} = W \cap \operatorname{Int} \mathcal{P}$, which contradicts the fact that W is a supporting hyperplane of $\mathcal{P}$ in Π. Thus, $q \in H \cap \operatorname{Int} \mathcal{K} \cap \operatorname{Int} \Pi^+$.

Next, since $p\vec{u} \subsetneq \mathcal{A}$, only the (unbounded) faces of $\mathcal{K}$ containing $\mathcal{A}$ intersect the half-line $p\vec{u}$. Let $\varepsilon_1 > 0$ be such that the ball $\mathcal{B}_n(p, \varepsilon_1)$ instersects none of the faces of $\mathcal{K}$ that do not contain $\mathcal{A}$. Consider now the line $\ell \subset H$ that contains $\mathcal{A}$ and the projection

$$\pi : H \to H \cap \Pi = W, \ x \mapsto (x + \vec{\ell}) \cap \Pi.$$

Since π is continuous and $\pi(p) = p$, there is $0 < \varepsilon_2 < \varepsilon_1$ such that $\pi(\mathcal{B}_n(p, \varepsilon_2)) \subset \mathcal{B}_n(p, \varepsilon_1)$. Fix $z \in \mathcal{B}_n(p, \varepsilon_2) \cap (\overline{pq} \setminus \{p\})$; we have $z \in \operatorname{Int} \mathcal{K}$. We claim that also $\pi(z) \in \operatorname{Int} \mathcal{K}$.

Indeed, let ℓ' be the line through z parallel to ℓ. Since $\vec{u} \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{\mathbf{0}\}$ and $\mathcal{K}$ is nondegenerated, there is $a \in \partial \mathcal{K}$ such that $a\vec{u} = \mathcal{K} \cap \ell'$; in fact, since $z \in \operatorname{Int} \mathcal{K}$, we have $a\vec{u} \setminus \{a\} \subset \operatorname{Int} \mathcal{K}$. Note also that a belongs only to faces of $\mathcal{K}$ that do not contain $\mathcal{A}$. Otherwise, a belongs to a face $\mathcal{E}$ of $\mathcal{K}$ containing $\mathcal{A}$, and since $\vec{u} \in \vec{\mathfrak{C}}(\mathcal{E})$, we deduce that $z \in a\vec{u} \subset \mathcal{E} \subset \partial \mathcal{K}$ but this contradicts the fact that $z \in \operatorname{Int} \mathcal{K}$. Thus, $a \notin \mathcal{B}_n(p, \varepsilon_1)$ because the ball $\mathcal{B}_n(p, \varepsilon_1)$ only intersects those faces of $\mathcal{K}$ containing $\mathcal{A}$. Since $\pi(z) \in \mathcal{B}_n(p, \varepsilon_1)$ and $z \in a\vec{u}$, we deduce that $\pi(z) \in a\vec{u} \setminus \{a\}$; otherwise, $\pi(z) \in a(-\vec{u})$ so that $a \in \overline{\pi(z)z} \subset \mathcal{B}_n(p, \varepsilon_1)$, a contradiction. Thus, $\pi(z) \in a\vec{u} \setminus \{a\} \subset \operatorname{Int} \mathcal{K}$.

Now, $W \cap \mathcal{P} \subset \partial \mathcal{P}$ because W is a supporting hyperplane of $\mathcal{P}$ in Π; hence

$$\pi(z) \in \pi(H) \cap \operatorname{Int} \mathcal{K} \subset H \cap \Pi \cap \operatorname{Int} \mathcal{K} = W \cap \operatorname{Int} \mathcal{P} = \varnothing,$$

a contradiction. Thus, $H \cap \operatorname{Int} \mathcal{K} = \varnothing$ and H is a supporting hyperplane of $\mathcal{K}$. □

The following result allows us to understand better nondegenerated unbounded convex polyhedra by comparing them with bounded convex polyhedra. This will provide later a more intuitive description of the recession cone for nondegenerated unbounded convex polyhedra. Along the proofs below we use implicitely the affine chart

$$\mathbb{RP}^n \setminus \{x_0 = 0\} \to \mathbb{R}^n, \quad (x_0 : \cdots : x_n) \mapsto (x_1/x_0, \ldots, x_n/x_0).$$

Lemma 1.6. *Let* $\mathcal{K} \subset \mathbb{R}^n \subset \mathbb{RP}^n$ *be a nondegenerated unbounded convex polyhedron in optimal* FU-*position with respect to* $\vec{\Pi} := \{x_n = 0\}$ *that does not intersect the hyperplane* $\Pi_0 := \{x_n = 0\}$. *Denote* $\widehat{\mathcal{K}} := \operatorname{Cl}_{\mathbb{RP}^n}(\mathcal{K})$ *and* $\mathcal{K}_\infty := \operatorname{Cl}_{\mathbb{RP}^n}(\mathcal{K}) \cap \mathsf{H}_\infty(\mathbb{R})$ *and consider the involutive homography*

$$\phi : \mathbb{RP}^n \to \mathbb{RP}^n, \ (x_0 : x_1 : \ldots : x_{n-1} : x_n) \mapsto (x_n : x_1 : \ldots : x_{n-1} : x_0),$$

whose restriction to $\mathbb{R}^n$ *is the rational map*

$$f := \phi|_{\mathbb{R}^n} : \mathbb{R}^n \dashrightarrow \mathbb{R}^n, \ (x_1, \ldots, x_n) \mapsto (y_1, \ldots, y_n) := \Big(\frac{x_1}{x_n}, \ldots, \frac{x_{n-1}}{x_n}, \frac{1}{x_n}\Big).$$

Then

(i) $\widehat{\mathcal{K}} = \mathcal{K} \sqcup \mathcal{K}_\infty$.

(ii) $\mathcal{K}' := \operatorname{Cl}(f(\mathcal{K})) = \phi(\widehat{\mathcal{K}}) \subset \mathbb{R}^n \equiv \{y_0 \neq 0\}$ *is a (nondegenerated) bounded convex polyhedron and one of its faces is* $\mathcal{E}' := \phi(\mathcal{K}_\infty)$.

(iii) *f is regular in $\mathcal{K}' \setminus \mathcal{E}'$ and satisfies the equality $f(\mathcal{K}' \setminus \mathcal{E}') = \mathcal{K}$. Moreover, the restriction map $f|_{\mathcal{K}'\setminus\mathcal{E}'} : \mathcal{K}'\setminus\mathcal{E}' \to \mathcal{K}$ is a biregular homeomorphism and $f(\operatorname{Int}\mathcal{K}') = \operatorname{Int}\mathcal{K}$.*

Proof. Statement (i) is clear because $\mathcal{K}$ is closed in $\mathbb{R}^n$. To prove (ii), we begin by proving that $\phi(\widehat{\mathcal{K}}) = \operatorname{Cl}(f(\mathcal{K}))$ is a bounded convex polyhedron. To that end, it is enough to check that $\phi(\widehat{\mathcal{K}}) \cap \mathsf{H}_\infty(\mathbb{R}) = \varnothing$, or equivalently, that $\phi(\widehat{\mathcal{K}}) \subset \mathbb{R}^n$.

Otherwise, since $\mathcal{K} \cap \Pi_0 = \varnothing$, there is $z := (0 : \vec{v}) \in \mathcal{K}_\infty \cap \operatorname{Cl}_{\mathbb{RP}^n}(\Pi_0)$; observe that $\vec{v} := (v_1, \dots, v_{n-1}, 0) \in \vec{\Pi}$. Since $\mathcal{K}$ is in optimal FU-position with respect to $\{x_n = 0\}$, there is a vertex $p := (p_1, \dots, p_n)$ of $\mathcal{K}$ such that $p_n < x_n$ for all $x := (x_1, \dots, x_n) \in \mathcal{K}\setminus\{p\}$. Let $\{z_k\}_k \subset \mathcal{K}\setminus\{p\}$ be a sequence that converges to z. For each $k \geq 1$, we choose a unitary vector $\vec{u}_k$ and $\lambda_k > 0$ such that $z_k = p + \lambda_k \vec{u}_k$. Moreover, since the unitary sphere $\mathbb{S}^{n-1}$ is compact, we may assume that the sequence $\{\vec{u}_k\}_k$ converges to a unitary vector $\vec{u}$; moreover $\lim_{k\to\infty} \lambda_k = +\infty$ because $z \in \mathcal{K}_\infty$. Observe that

$$z_k \equiv (1 : z_k) = \Big(\frac{1}{\lambda_k} : \frac{p}{\lambda_k} + \vec{u}_k\Big)$$

tends to $(0 : \vec{u})$ when $k \to +\infty$. Thus, we may assume that $\vec{u} = \vec{v} = (v_1, \dots, v_{n-1}, 0)$ and $z := (0 : \vec{u})$. Let us see now that the half-line $p\vec{u} \subset \mathcal{K}$. Indeed, let $x \in p\vec{u}$; since $\vec{u}$ is unitary, $x = p + \|x - p\|\vec{u}$. Let $k_0 \geq 1$ be such that for each $k \geq k_0$ it holds that $\lambda_k > \|x - p\|$; hence,

$$y_k = p + \|x - p\|\vec{u}_k \in \overline{pz_k} = \{p + t\vec{u}_k : \ 0 \leq t \leq \lambda_k\} \subset \mathcal{K}.$$

Since $\mathcal{K}$ is closed and

$$\lim_{k\to\infty} y_k = p + \|x - p\|\vec{u} = x$$

we deduce that $x \in \mathcal{K}$. Thus,

$$\{(p_1 + tv_1, \dots, p_{n-1} + tv_{n-1}, p_n) : \ t \geq 0\} = p\vec{u} \subset \mathcal{K},$$

against the fact that $p_n < x_n$ for all $x := (x_1, \dots, x_n) \in \mathcal{K} \setminus \{p\}$.

Next, since $\mathcal{K}$ is unbounded and connected, $\{y_n = 0\}$ is a supporting hyperplane of $\phi(\widehat{\mathcal{K}})$; hence, $\mathcal{E}' := \phi(\mathcal{K}_\infty) = \phi(\widehat{\mathcal{K}}) \cap \{y_n = 0\}$ is a face of $\phi(\widehat{\mathcal{K}})$.

(iii) Since

$$\mathcal{K}' \setminus \mathcal{E}' = \mathcal{K}' \setminus \{y_n = 0\} = \operatorname{Cl}(f(\mathcal{K})) \setminus \{y_n = 0\} = f(\mathcal{K}),$$

we have $\mathcal{K} = f(f(\mathcal{K})) = f(\mathcal{K}' \setminus \mathcal{E}')$. Moreover, $f|_{\mathcal{K}'\setminus\mathcal{E}'} : \mathcal{K}' \setminus \mathcal{E}' \to \mathcal{K}$ is a biregular homeomorphism whose inverse is the restriction map $f|_{\mathcal{K}} : \mathcal{K} \to \mathcal{K}' \setminus \mathcal{E}'$. Moreover, since $f|_{\mathcal{K}'\setminus\mathcal{E}'}$ is a diffeomorfism, it transforms $\operatorname{Int}\mathcal{K}' = \operatorname{Int}(\mathcal{K}' \setminus \mathcal{E}')$ onto $\operatorname{Int}\mathcal{K}$. □

1.4. **Structure of the recession cone of a nondegenerated convex polyhedron.** The following result provides a clear description of the structure of the recession cone of a nondegenerated unbounded convex polyhedron.

Proposition 1.7. *Let $\mathcal{K} \subset \mathbb{R}^n \subset \mathbb{RP}^n$ be a nondegenerated unbounded convex polyhedron in* FU*-position with respect to the hyperplane $\vec{\Pi} := \{x_n = 0\}$ and which does not intersect the hyperplane $\Pi_0 := \{x_n = 0\}$. Denote $\widehat{\mathcal{K}} := \operatorname{Cl}_{\mathbb{RP}^n}(\mathcal{K})$ and $\mathcal{K}_\infty := \operatorname{Cl}_{\mathbb{RP}^n}(\mathcal{K}) \cap \mathsf{H}_\infty(\mathbb{R})$. Consider the involutive homography*

$$\phi : \mathbb{RP}^n \to \mathbb{RP}^n, \ (x_0 : x_1 : \ldots : x_{n-1} : x_n) \mapsto (x_n : x_1 : \ldots : x_{n-1} : x_0).$$

Fix $p \in \mathcal{K}$. Then

(i) *$\phi(\mathfrak{C}_p(\mathcal{K})) \cup \phi(\mathcal{K}_\infty)$ is the closed cone $\mathcal{C}_{\phi(p)}$ of base $\phi(\mathcal{K}_\infty)$ and vertex $\phi(p)$; hence, $\phi(\mathfrak{C}_p(\mathcal{K})) = \mathcal{C}_{\phi(p)} \setminus \phi(\mathcal{K}_\infty)$. In particular, $\mathfrak{C}_p(\mathcal{K})$ is an unbounded convex polyhedron for each $p \in \mathcal{K}$.*

(ii) The projection $\pi_p : \mathbb{RP}^n \setminus \{p\} \to \mathsf{H}_\infty(\mathbb{R})$ of center $\{p\}$ and base $\mathsf{H}_\infty(\mathbb{R})$, defined by $\pi_p(x_0 : x) = (0 : x - px_0)$, where $x := (x_1, \dots, x_n)$ satisfies
- $\pi_p(\mathfrak{C}_p(\mathcal{K}) \setminus \{p\}) = \mathcal{K}_\infty$,
- $\mathfrak{C}_p(\mathcal{K}) = (\pi_p^{-1}(\mathcal{K}_\infty) \cap \{x_n - p_n > 0\}) \cup \{p\}$; and
- $\pi_p(\mathrm{Int}(\mathfrak{C}_p(\mathcal{K}))) = \mathrm{Int}(\mathcal{K}_\infty)$ and $\pi_p(\partial\mathfrak{C}_p(\mathcal{K}) \setminus \{p\}) = \partial\mathcal{K}_\infty$.

Proof. First, let us see that $\pi_p(p + \vec{\mathfrak{C}}(\mathcal{K})) = \mathcal{K}_\infty$ for each $p \in \mathcal{K}$. Let $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{\mathbf{0}\}$ and for $k \in \mathbb{N}$ define $z_k := p + k\vec{v} \in p\vec{v} \subset \mathcal{K}$. The point $q := (0 : \vec{v}) \in \mathsf{H}_\infty(\mathbb{R})$ is the limit in $\mathbb{RP}^n$ of the sequence $\{z_k\}_k$; hence,

$$\pi_p(p + \vec{v}) = (0 : \vec{v}) = q \in \widehat{\mathcal{K}} \cap \mathsf{H}_\infty(\mathbb{R}) = \mathcal{K}_\infty$$

and so $\pi_p(\mathfrak{C}(\mathcal{K}) \setminus \{p\}) \subset \mathcal{K}_\infty$. Conversely, if $q := (0 : \vec{v}) \in \mathcal{K}_\infty$, the segment $\overline{pq} \subset \widehat{\mathcal{K}}$; hence, $p\vec{v} \subset \mathcal{K}$ and so $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})$ and $q = \pi_p(p + \vec{v}) \in \pi_p(\mathfrak{C}(\mathcal{K}) \setminus \{p\})$.

(i) Since $\mathfrak{C}_p(\mathcal{K}) = p + \vec{\mathfrak{C}}(\mathcal{K}) = \bigcup_{\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})} p\vec{v}$, we have

$$\phi(\mathfrak{C}_p(\mathcal{K})) = \bigcup_{\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})} \phi(p\vec{v}) = \bigcup_{\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})} \overline{\phi(p)\phi(0 : \vec{v})} \setminus \{\phi(0 : \vec{v})\} = \mathcal{C}_{\phi(p)} \setminus \phi(\mathcal{K}_\infty).$$

Since $\mathcal{C}_{\phi(p)}$ is a convex polyhedron, because it is the cone of vertex $\phi(p)$ and base $\phi(\mathcal{K}_\infty)$, we deduce that $\mathfrak{C}_p(\mathcal{K}) = \phi(\mathcal{C}_{\phi(p)}) \setminus \mathcal{K}_\infty$ is an unbounded convex polyhedron for each $p \in \mathcal{K}$.

(ii) We have already proved that $\pi_p(\mathfrak{C}_p(\mathcal{K})) = \mathcal{K}_\infty$. As a consequence of this and the fact that $\mathcal{K}$ is in FU-position with respect to $\{x_n = 0\}$, we deduce that

$$\mathfrak{C}_p(\mathcal{K}) \subset (\pi_p^{-1}(\mathcal{K}_\infty) \cap \{x_n - p_n > 0\}) \cup \{p\}$$

To prove the converse inclusion, let $\vec{v} := (v_1, \dots, v_n) \in \mathbb{R}^n$ be such that $\pi_p(p + \vec{v}) \in \mathcal{K}_\infty$ and $v_n > 0$. Thus, there is $\vec{w} := (w_1, \dots, w_n) \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{\mathbf{0}\}$ such that $\pi_p(p + \vec{w}) = \pi_p(p + \vec{v})$, that is, there is $\lambda \neq 0$ such that $\vec{v} = \lambda\vec{w}$. Since $\vec{\mathfrak{C}}(\mathcal{K}) \subset \{x_n > 0\}$, we have $w_n > 0$ and so, $\lambda > 0$; hence, $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})$.

It only remains to prove that $\pi_p(\mathrm{Int}(\mathfrak{C}_p(\mathcal{K}))) = \mathrm{Int}(\mathcal{K}_\infty)$ and $\pi_p(\partial\mathfrak{C}_p(\mathcal{K}) \setminus \{p\}) = \partial\mathcal{K}_\infty$. Since $\pi_p(\mathfrak{C}_p(\mathcal{K}) \setminus \{p\}) = \mathcal{K}_\infty$, it is enough to check that $\pi_p(\mathrm{Int}(\mathfrak{C}_p(\mathcal{K}))) \subset \mathrm{Int}(\mathcal{K}_\infty)$ and $\pi_p(\partial\mathfrak{C}_p(\mathcal{K}) \setminus \{p\}) \subset \partial\mathcal{K}_\infty$. We leave the concrete details to the reader. □

As a consequence of the previous result, it is not difficult to generalize, for nondegenerated unbounded convex polyhedra, the fact that a bounded convex polyhedron coincides with the convex hull of the set of its vertices, see §1.1.3. We leave the concrete details of the proof to the reader (see also [Z, Ch.1]).

Corollary 1.8. *Let $\mathcal{K} \subset \mathbb{R}^n$ be a nondegenerated unbounded convex polyhedron, let $\mathfrak{V} := \{p_1, \dots, p_r\}$ be the collection of its vertices, let $\mathcal{A}_1, \dots, \mathcal{A}_s$ be its unbounded edges and for each of them let $\vec{v}_j \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{0\}$ be such that $\mathcal{A}_j$ is parallel to $\vec{v}_j$. Then*

(i) $\mathcal{K} = \{\sum_{i=1}^r \mu_i p_i + \sum_{j=1}^s \lambda_j \vec{v}_j : \ \lambda_j, \mu_i \geq 0, \ \sum_{i=1}^r \mu_i = 1\}$

(ii) $\vec{\mathfrak{C}}(\mathcal{K}) = \{\sum_{j=1}^s \lambda_j \vec{v}_j : \ \lambda_1, \dots, \lambda_s \geq 0\}$.

(iii) $\mathcal{K} = \mathcal{K}_0 + \vec{\mathfrak{C}}(\mathcal{K})$ *where* $\mathcal{K}_0 := \{\sum_{i=1}^r \mu_i p_i : \ \sum_{i=1}^r \mu_i = 1, \mu_i \geq 0\}$ *is a bounded convex polyhedron.*

Next, we show that if the recession cone $\vec{\mathfrak{C}}(\mathcal{K})$ of a convex polyhedron $\mathcal{K} \subset \mathbb{R}^n$ has maximal dimension, then any line parallel to a vector of $\operatorname{Int}\vec{\mathfrak{C}}(\mathcal{K})$ intersects $\operatorname{Int}\mathcal{K}$ in a half-line.

Lemma 1.9. *Let $\mathcal{K} \subsetneq \mathbb{R}^n$ be an n-dimensional unbounded convex polyhedron such that $\dim(\vec{\mathfrak{C}}(\mathcal{K})) = n$ and let $\vec{v} \in \operatorname{Int}\vec{\mathfrak{C}}(\mathcal{K})$. Then:*

(i) *For each $p \in \mathbb{R}^n$ the intersection $p\vec{v} \cap \mathcal{K}$ is a half-line $p_1\vec{v} \subset \mathcal{K}$.*

(ii) *If $\mathcal{K}$ is nondegenerated and it is in* FU*-position with respect to a hyperplane $\vec{\Pi}$ and $\mathfrak{F} \subset \mathbb{R}^n$ is a finite set, there is a hyperplane Π' parallel to $\vec{\Pi}$ such that the intersection $p\vec{v} \cap (\mathcal{K} \cap \Pi')$ is a singleton for each $p \in \mathfrak{F}$.*

Proof. (i) If $p \in \mathcal{K}$, the intersection $p\vec{v} \cap \mathcal{K} = p\vec{v}$ is a half-line and there is nothing to be proved; hence, assume that $p \notin \mathcal{K}$. If $n = 1$, the polyhedron $\mathcal{K}$ is a half-line and the statement holds. Suppose first that $n \geq 2$ and $p \in \mathbb{R}^n \setminus \mathcal{K}$; we argue by way of contradiction. Since $n \geq 2$ there is a point $y \in \mathcal{K}$ not belonging to the line ℓ generated by the points p and $p + \vec{v}$. Since $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})$, we have $y\vec{v} \subset \mathcal{K}$ and, since $y \notin \ell$, there is a unique plane $\pi \subset \mathbb{R}^n$ through the points y, p and $p + \vec{v}$.

Since $\vec{v} \in \operatorname{Int}(\vec{\mathfrak{C}}(\mathcal{K}))$, there is $\varepsilon > 0$ such that the vector $\vec{w} := \vec{v} + \varepsilon(p - y) \in \vec{\mathfrak{C}}(\mathcal{K})$; hence, $y\vec{w} \subset \mathcal{K}$. Observe that the half-lines $y\vec{w}$ and $p\vec{v}$ meet at the point

$$z := p + \frac{1}{\varepsilon}\vec{v} = y + \frac{1}{\varepsilon}(\vec{v} + \varepsilon(p - y)) = y + \frac{1}{\varepsilon}\vec{w} \in \mathcal{K}.$$

Since $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K})$, the half-line $z\vec{v} \subset \mathcal{K}$ and so the convex set $p\vec{v} \cap \mathcal{K}$ contains the half-line $z\vec{v}$; hence, it is also a half-line.

(ii) Write $\vec{\Pi} := \{\vec{h} = 0\}$ and let $\Pi := \{h = 0\}$ be a parallel hyperplane to $\vec{\Pi}$ that meet all the unbounded edges $\mathcal{A}_1, \ldots, \mathcal{A}_s$ of $\mathcal{K}$ and such all the vertices of $\mathcal{K}$ are contained in $\operatorname{Int}\Pi^-$. Write $\mathcal{A}_i := q_i\vec{v}_i$ where q_i is a vertex of $\mathcal{K}$, the vector $\vec{v}_i \in \vec{\mathfrak{C}}(\mathcal{K})$ and $b_i := q_i + \vec{v}_i \in \Pi$ for $i = 1, \ldots, s$. Since $0 = h(b_i) = h(q_i) + \vec{h}(\vec{v}_i)$ and $h(q_i) < 0$, we deduce that $\vec{h}(\vec{v}_i) = -h(q_i) > 0$. By Corollary 1.8, there are $\lambda_i \geq 0$ not all zero such that $\vec{v} = \sum_{i=1}^s \lambda_i\vec{v}_i$; hence,

$$\vec{h}(\vec{v}) = \vec{h}\Big(\sum_{i=1}^s \lambda_i\vec{v}_i\Big) = \sum_{i=1}^s \lambda_i\vec{h}(\vec{v}_i) > 0.$$

By statement (i), for each $p \in \mathfrak{F}$ there is a point $p' \in \mathcal{K}$ such that $p\vec{v} \cap \mathcal{K} = p'\vec{v}$. We choose $\Pi' = \{h' = 0\}$ parallel to $\vec{\Pi}$ such that $p' \in \operatorname{Int}\Pi'^-$ for each $p \in \mathfrak{F}$. We claim that $p\vec{v} \cap (\mathcal{K} \cap \Pi') \neq \varnothing$ for each $p \in \mathfrak{F}$.

Indeed, since $\vec{h}' = \vec{h}$, we have $\vec{h}'(\vec{v}) > 0$. Fix $p \in \mathfrak{F}$; since $h'(p') < 0$, there is $t > 0$ such that $h'(p' + t\vec{v}) = 0$ and so $p' + t\vec{v} \in p'\vec{v} \cap \Pi' = p\vec{v} \cap (\mathcal{K} \cap \Pi')$. Moreover, since $p'\vec{v}$ is a half-line and $p' \notin \Pi'$, the intersection $p\vec{v} \cap (\mathcal{K} \cap \Pi')$ is a singleton. □

2. On sections and projections of convex polyhedra

The purpose of this section is to understand how are the generic sections of a nondegenerated unbounded convex polyhedron and to decide when the projection of a convex polyhedron coincides off a compact set with one of its sections.

Lemma 2.1. *Let $\mathcal{K} \subset \mathbb{R}^n$ be an n-dimensional nondegenerated unbounded convex polyhedron and let $\vec{\Pi} := \{\vec{h} = 0\}$ be an (oriented vectorial) hyperplane. Let $\mathcal{A}_1, \ldots, \mathcal{A}_s$ be the unbounded edges of $\mathcal{K}$, ordered in such a way that the first k (where $0 \leq k \leq s$) are* *nonparallel to* $\vec{\Pi}$ *and the remaining* $s-k$ *are parallel to* $\vec{\Pi}$*; for each* $i=1,\dots,s$ *write* $\mathcal{A}_i := q_i\vec{v}_i$ *where* q_i *is a vertex of* $\mathcal{K}$ *and* $\vec{v}_i \in \vec{\mathfrak{C}}(\mathcal{K})$ *for* $i=1,\dots,s$*. Let* Π *be a hyperplane parallel to* $\vec{\Pi}$ *that meets* $\mathcal{A}_1,\dots,\mathcal{A}_k$ *and such all the vertices of* $\mathcal{K}$ *and the unbounded edges* $\mathcal{A}_{k+1},\dots,\mathcal{A}_s$ *are contained in* $\operatorname{Int}\Pi^-$*. Moreover, let* $\Pi' \subset \Pi^+$ *be a hyperplane parallel to* $\vec{\Pi}$*, let* $\mathfrak{H} := \{H_1,\dots,H_m\}$ *be the minimal presentation of* $\mathcal{K}$*, and let* $\mathfrak{H}_1 := \{H_{i_1},\dots,H_{i_r}\}$ *be the collection of hyperplanes in* $\mathfrak{H}$ *containing any of the edges* $\mathcal{A}_1,\dots,\mathcal{A}_k$*. Then we have:*

- (i) *The intersection* $\mathcal{A}_i \cap \Pi'^+$ *is a half-line whose boundary is* $\mathcal{A}_i \cap \Pi'$ *for* $i=1,\dots,k$*.*
- (ii) *The polyhedron* $\mathcal{P}' := \mathcal{K}\cap\Pi' \neq \varnothing$ *if and only if* $k \geq 1$*.*
- (iii) *If* $\mathcal{P}' \neq \varnothing$*, then* $\mathcal{P}'$ *is a convex polyhedron whose vertices are the intersections of* Π' *with the unbounded edges of* $\mathcal{K}$ *(nonparallel to* $\vec{\Pi}$*) and whose recession cone is* $\vec{\mathfrak{C}}(\mathcal{P}') = \{\sum_{j=k+1}^s \lambda_j\vec{v}_j :\ \lambda_j \geq 0\}$*.*
- (iv) *The recession cone* $\vec{\mathfrak{C}}(\mathcal{K}\cap\Pi'^-) = \mathcal{C} := \{\sum_{j=k+1}^s \lambda_j\vec{v}_j :\ \lambda_j \geq 0\}$*, and the edges* $\mathcal{A}_{k+1},\dots,\mathcal{A}_s$*, together with all the bounded faces of* $\mathcal{K}$*, are contained in* $\mathcal{K}\cap\operatorname{Int}\Pi'^-$*.*
- (v) *The hyperplane* Π' *meets all the unbounded faces of* $\mathcal{K}$ *that contain one of the edges* $\mathcal{A}_1,\dots,\mathcal{A}_k$ *and just these faces.*
- (vi) *If* $\mathcal{P}' \neq \varnothing$*, the minimal presentation of* $\mathcal{P}'$ *in* Π' *is* $\{H_{i_1}\cap\Pi',\dots,H_{i_t}\cap\Pi'\}$*. In particular, the facets of* $\mathcal{P}'$ *are* $H_{i_\ell}\cap\mathcal{P}'$ *for* $\ell=1,\dots,t$*,*
- (vii) *If* $\mathcal{P}' \neq \varnothing$*, then* $\mathcal{K}\cap\Pi'^+ = \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{K}) = \mathcal{P}' + \{\sum_{j=1}^k \lambda_j\vec{v}_j :\ \lambda_j \geq 0\}$*.*

Proof. (i) By hypothesis the vertex $q_j \in \operatorname{Int}\Pi^-$, that is, $h(q_j) < 0$ and since $\Pi\cap\mathcal{A}_j \neq \varnothing$, we deduce that $\vec{h}(\vec{v}_i) > 0$, and so $\mathcal{A}_j \cap \Pi'^+$ is a half-line, whose boundary is $\Pi'\cap\mathcal{A}_j$.

(ii) If $k \geq 1$, then $\mathcal{P}' := \mathcal{K}\cap\Pi' \supset \mathcal{A}_1\cap\Pi' \neq \varnothing$. Conversely, if $k=0$, denote $p_1,\dots,p_r$ the vertices of $\mathcal{K}$. By Corollary 1.8, we have

$$\mathcal{K} = \Big\{\sum_{i=1}^r \mu_i p_i + \sum_{j=1}^s \lambda_j\vec{v}_j :\ \mu_i,\lambda_j \geq 0,\ \sum_{i=1}^r \mu_i = 1\Big\}$$

and this last set is contained in $\operatorname{Int}\Pi'^-$ because $p_1,\dots,p_r \in \operatorname{Int}\Pi'^-$ and the edges $\mathcal{A}_j := q_j\vec{v}_j \subset \operatorname{Int}\Pi^-$ for $j=1,\dots,s$. Thus, $\mathcal{P}' = \mathcal{K}\cap\Pi' = \varnothing$.

(iii) and (iv) Assume that $\mathcal{P}'$ is nonempty; in particular, it is a convex polyhedron. Let p be a vertex of $\mathcal{P}'$; notice that $p \in \partial\mathcal{K}$. Choose the face $\mathcal{E}$ of $\mathcal{K}$ of least dimension between those containing p; then, $p \in \operatorname{Int}\mathcal{E}$. Since all the vertices of $\mathcal{K}$ are contained in $\operatorname{Int}\Pi^-$ and $\Pi' \subset \Pi^+$, we deduce that the dimension of $\mathcal{E}$ is ≥ 1. Let W be the affine subspace generated by $\mathcal{E}$ and let $\varepsilon > 0$ be such that $\mathcal{B}_n(p,\varepsilon)\cap W \subset \mathcal{E}$. Since Π' is a hyperplane we deduce that $\dim W - 1 \leq \dim(W\cap\Pi') \leq \dim W$. Since $\mathcal{B}_n(p,\varepsilon)\cap W \subset \mathcal{E}$,

$$\mathcal{B}_n(p,\varepsilon)\cap W\cap\Pi' \subset \mathcal{E}\cap\Pi' = \mathcal{E}\cap\mathcal{P}' \subset \mathcal{P}';$$

hence, since p is a vertex of $\mathcal{P}'$ we have $W\cap\Pi' = \{p\}$ (because otherwise we would have a face of $\mathcal{P}'$ crossing the vertex), and so $0 \leq \dim W - 1 \leq \dim(W\cap\Pi') = 0$. Thus, $\mathcal{E} := \mathcal{A}_j$ is an (unbounded) edge of $\mathcal{K}$ that meets Π' in a unique point. This means that $\mathcal{A}_j$ is nonparallel to $\vec{\Pi}$.

Conversely, if $\mathcal{A}_j$ is an unbounded edge of $\mathcal{K}$ nonparallel to $\vec{\Pi}$, then the intersection $\mathcal{A}_j\cap\Pi' = \{p\}$. On the other hand, let H be a supporting hyperplane of $\mathcal{K}$ such that $\mathcal{K}\cap H = \mathcal{A}_j$. Thus, the intersection $\mathcal{P}'\cap H = \mathcal{A}_j\cap\Pi = \{p\}$ and p is a vertex of $\mathcal{P}'$. This completes the part concerning the determination of the vertices of $\mathcal{P}'$.

Observe now that if $\mathcal{E}$ is a bounded face of $\mathcal{K}$, then $\mathcal{E}$ coincides with the convex hull of its vertices, which are also vertices of $\mathcal{K}$, and they are contained in the open half-space $\operatorname{Int}\Pi^- \subset \operatorname{Int}\Pi'^-$; hence, $\mathcal{E} \subset \Pi'^-$. Moreover, by hypothesis $\mathcal{A}_j \subset \Pi'^-$ for $j = k+1, \dots, s$.

To end the proof of statements (iii) and (iv) it only remains to prove the equalities

$$\vec{\mathfrak{C}}(\mathcal{P}') = \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^-) = \mathcal{C} := \Big\{ \sum_{j=k+1}^{s} \lambda_j \vec{v}_j : \ \lambda_j \geq 0 \Big\}.$$

Indeed, let $h := \vec{h} + a \in \mathbb{R}[\mathrm{x}]$ be a degree one equation of Π, such that $\Pi^+ = \{h \geq 0\}$; in particular $\Pi' = \{h' := \vec{h} + b = 0\}$ for some $b \geq 0$. Using (i), one sees that

$$\vec{h}(\vec{v}_i) \begin{cases} > 0 & \text{para } i = 1, \dots, k, \\ = 0 & \text{para } i = k+1, \dots, s. \end{cases} \tag{2.1}$$

Let $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^-)$; then $\vec{v} \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{0\}$ and there is $p \in \mathcal{K} \cap \Pi'^-$ such that $p\vec{v} \subset \mathcal{K} \cap \Pi'^-$; hence $\vec{h}(\vec{v}) \leq 0$. By Corollary 1.8, $\vec{\mathfrak{C}}(\mathcal{K}) = \{\sum_{j=1}^{s} \lambda_j \vec{v}_j : \ \lambda_j \geq 0\}$. In particular, there are $\zeta_j \geq 0$ such that $\vec{v} = \sum_{j=1}^{s} \zeta_j \vec{v}_j$. Thus, since $\vec{v}_j \in \vec{\Pi}$ and $\vec{h}(\vec{v}_j) = 0$ for $j = k+1, \dots, s$, we deduce that

$$0 \geq \vec{h}(\vec{v}) = \vec{h}\Big(\sum_{j=1}^{s} \zeta_j \vec{v}_j\Big) = \sum_{j=1}^{s} \zeta_j \vec{h}(\vec{v}_j) = \sum_{j=1}^{k} \zeta_j \vec{h}(\vec{v}_j).$$

But $\vec{h}(\vec{v}_j) > 0$ for $j = 1, \dots, k$ and each $\zeta_j \geq 0$; hence, $\mu_j = 0$ for $j = 1, \dots, k$, and this means that $\vec{v} \in \mathcal{C}$. Thus,

$$\vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^-) \subset \mathcal{C} \subset \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{\Pi} = \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi') = \vec{\mathfrak{C}}(\mathcal{P}') \subset \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^-),$$

and so the desired equalities are true. Observe that if $\mathcal{P}' = \varnothing$, one proceeds analogously, using that

$$\mathcal{C} \subset \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{\mathfrak{C}}(\Pi'^-) = \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^-).$$

(v) Let $\mathcal{E}$ be a face of $\mathcal{K}$. If $\mathcal{E}$ is bounded, we have already seen that $\mathcal{E} \subset \operatorname{Int}\Pi'^-$ and so it does not intersect Π'. Thus, assume that $\mathcal{E}$ is unbounded; since $\mathcal{K}$ is nondegenerated, then $\mathcal{E}$ is nondegenerated (because otherwise it would contain a line and this would imply that $\mathcal{K}$ is degenerate, see [FGU, §2.2] for further details) and in particular $\mathcal{E}$ has some unbounded edges, which are also (unbounded) edges of $\mathcal{K}$. If any of the previous edges is $\mathcal{A}_i$ for $i = 1, \dots, k$, then $\mathcal{E} \cap \Pi' \supset \mathcal{A}_i \cap \Pi' \neq \varnothing$. Suppose now that all the unbounded edges of $\mathcal{E}$ are between $\mathcal{A}_{k+1}, \dots, \mathcal{A}_s$ and let $p_1, \dots, p_r$ be the vertices of $\mathcal{K}$. Obviously, the vertices of $\mathcal{E}$ constitute a subset of the vertices of $\mathcal{K}$. Then, by Corollary 1.8 we deduce

$$\mathcal{E} \subset \Big\{ \sum_{i=1}^{r} \mu_i p_i + \sum_{j=k+1}^{s} \lambda_j \vec{v}_j : \ \mu_i, \lambda_j \geq 0, \ \sum_{i=1}^{r} \mu_i = 1 \Big\} \subset \operatorname{Int}\Pi'^-,$$

because $p_1, \dots, p_r \in \operatorname{Int}\Pi'^-$ and $\vec{v}_{k+1}, \dots, \vec{v}_s \in \vec{\Pi}$. Thus, $\mathcal{E} \cap \Pi' = \varnothing$.

(vi) First, we claim that the hyperplane Π' meets the interior of $\mathcal{K}$. Otherwise $\varnothing \neq \mathcal{P}' \subset \partial\mathcal{K}$ and we deduce that Π' is a supporting hyperplane of $\mathcal{K}$. Thus, $\mathcal{P}'$ is a face of $\mathcal{K}$; moreover, by (iii) above, $\vec{\mathfrak{C}}(\mathcal{P}') = \{\sum_{j=k+1}^{s} \lambda_j \vec{v}_j : \ \lambda_j \geq 0\}$. But this is against the fact that Π' only meets the faces of $\mathcal{K}$ that contain one of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$. Thus, Π' meets the interior of $\mathcal{K}$ and so, the dimension of $\mathcal{P}'$ is $n-1$.

Moreover, by [Be2, 12.1.7] the facets of $\mathcal{P}'$ are those intersections of facets of $\mathcal{K}$ with Π' that have dimension $n-2$. Recall that the facets $\mathcal{F}$ of $\mathcal{K}$ which do not contain one of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$ do not meet Π'; hence, these facets of $\mathcal{K}$ do not provide facets of

$\mathcal{P}'$. On the other hand, let us see that for each $\ell = 1, \dots, t$ the intersection of Π' with $\mathcal{F}_{i_\ell} := H_{i_\ell} \cap \mathcal{K}$ has dimension $n-2$. To that end, it is enough to prove that Π' meets the interior of $\mathcal{F}_{i_\ell}$. Otherwise, $\mathcal{F}_{i_\ell} \cap \Pi' \subset \partial\mathcal{F}_{i_\ell} \subset \partial\mathcal{K}$ and since Π' meets and only meets the faces of $\mathcal{K}$ that contain one of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$, we deduce that $\mathcal{F}_{i_\ell} \cap \Pi'$ is a face of $\mathcal{K}$ (and so of $\mathcal{F}_{i_\ell}$) that contains one of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$. But this is impossible because $\mathcal{F}_{i_\ell} \cap \Pi' \subset \mathcal{K} \cap \Pi' = \mathcal{P}'$ and $\vec{\mathfrak{C}}(\mathcal{P}') = \{\sum_{j=k+1}^s \lambda_j \vec{v}_j : \ \lambda_j \geq 0\}$. Thus, $H_{i_\ell} \cap \mathcal{P}' = \mathcal{F}_{i_\ell} \cap \Pi'$ has dimension $n-2$ and it is a facet of $\mathcal{P}'$ for $\ell = 1, \dots, t$.

Let us see that if $j \neq \ell$, then the facets $H_{i_\ell} \cap \mathcal{P}'$ and $H_{i_{\ell'}} \cap \mathcal{P}'$ are different. Otherwise, the intersection $H_{i_\ell} \cap \mathcal{P}' = H_{i_\ell} \cap H_{i_{\ell'}} \cap \mathcal{P}'$ has dimension $n-2$ and is contained in the affine subspace $H_{i_\ell} \cap H_{i_{\ell'}}$, which also has dimension $n-2$. In this way, $H_{i_\ell} \cap \mathcal{P}' \subset \Pi'$ generates $H_{i_\ell} \cap H_{i_{\ell'}}$ and so $H_{i_\ell} \cap H_{i_{\ell'}} \subset \Pi'$. Since Π' only meets the faces of $\mathcal{K}$ that contain one of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$, we deduce that $H_{i_\ell} \cap H_{i_{\ell'}} \cap \mathcal{K}$ contains one of those edges. But since

$$H_{i_\ell} \cap H_{i_{\ell'}} \cap \mathcal{K} = H_{i_\ell} \cap H_{i_{\ell'}} \cap \mathcal{P}'$$

is a facet of $\mathcal{P}'$ and $\vec{\mathfrak{C}}(\mathcal{P}') = \{\sum_{j=k+1}^s \lambda_j \vec{v}_j : \ \lambda_j \geq 0\}$, we arrive at a contradiction. Thus, the (different) facets of $\mathcal{P}'$ are $H_{i_\ell} \cap \mathcal{P}'$ for $\ell = 1, \dots, t$. Moreover, since $\mathcal{F}_{i_\ell}$ generates the affine subspace $H_{i_\ell} \cap \Pi'$, we deduce that $\{H_{i_1} \cap \Pi', \dots, H_{i_t} \cap \Pi'\}$ is the minimal presentation of the convex polyhedron $\mathcal{P}'$ in Π'.

(vii) If $\mathcal{P}' \neq \varnothing$, the vertices of the nondegenerated convex polyhedron $\mathcal{K} \cap \Pi'^+$ are the intersections of Π' with the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$ of $\mathcal{K}$ (see (iii)). Thus, the convex hull of the set constituted by those vertices is contained in $\mathcal{P}'$ and by Corollary 1.8 (iii), $\mathcal{K} \cap \Pi'^+ = \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^+)$. Moreover, by equation (2.1) and Corollary 1.8, we deduce

$$\vec{\mathfrak{C}}(\mathcal{K}) = \Big\{ \sum_{j=1}^s \lambda_j \vec{v}_j : \ \lambda_j \geq 0 \Big\} \subset \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^+) \subset \vec{\mathfrak{C}}(\mathcal{K}).$$

Moreover, since $\vec{\mathfrak{C}}(\mathcal{P}') = \{\sum_{j=k+1}^s \lambda_j \vec{v}_j : \ \lambda_j \geq 0\}$ we deduce that

$$\begin{aligned} \mathcal{K} \cap \Pi'^+ &= \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi'^+) = \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{K}) \\ &= \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{P}') + \Big\{ \sum_{j=1}^k \lambda_j \vec{v}_j : \ \lambda_j \geq 0 \Big\} = \mathcal{P}' + \Big\{ \sum_{j=1}^k \lambda_j \vec{v}_j : \ \lambda_j \geq 0 \Big\} \end{aligned}$$

and we are done. □

Remark 2.2. If $k = s$ in the previous Lemma 2.1, then $\mathcal{K}$ is in FU-position with respect to $\vec{\Pi}$ and it holds:

(i) The intersection $\mathcal{A}_i \cap \Pi'^+$ is a half-line whose boundary is $\mathcal{A}_i \cap \Pi'$ for $i = 1, \dots, s$.
(ii) The convex polyhedron $\mathcal{P}' := \mathcal{K} \cap \Pi' \neq \varnothing$ is bounded and its vertices are the intersections of Π' with the unbounded edges of $\mathcal{K}$,
(iii) The polyhedron $\mathcal{K} \cap \Pi'^-$ is bounded and all the bounded faces of $\mathcal{K}$ are contained in $\mathcal{K} \cap \operatorname{Int} \Pi'^-$.
(iv) Π' meet all the unbounded faces of $\mathcal{K}$ and just these faces.
(v) The minimal presentation of $\mathcal{P}'$ is $\{H_{i_1} \cap \Pi', \dots, H_{i_r} \cap \Pi'\}$. In particular, the facets of $\mathcal{P}'$ are $H_{i_j} \cap \mathcal{P}'$ for $j = 1, \dots, r$,
(vi) $\mathcal{K} \cap \Pi'^+ = \mathcal{P}' + \vec{\mathfrak{C}}(\mathcal{K})$.

The following result concerning projections of convex polyhedra is the clue to characterize when convex polyhedra can be placed in first or second trimming position. Recall that

$$\pi_n:\mathbb{R}^n\to\mathbb{R}^{n-1}\times\{0\},\ x:=(x_1,\dots,x_n)\mapsto(x',0):=(x_1,\dots,x_{n-1},0)$$

and we denote by $\vec{\ell}_n$ the vectorial line generated by the vector $\vec{e}_n:=(0,\dots,0,1)$.

Proposition 2.3 (Projections of convex polyhedra)**.** *Let $\mathcal{K}\subset\mathbb{R}^n$ be a nondegenerated unbounded convex polyhedron. Then the set*

$$\Lambda:=\{a\in\mathbb{R}^{n-1}:\ \mathcal{I}_a:=\pi_n^{-1}(a,0)\cap\mathcal{K}\neq\varnothing,\ (a,0)\not\in\mathcal{I}_a\}$$

is bounded if and only if the following statements hold:

- (i) *The projection $\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))=\vec{\mathfrak{C}}(\mathcal{K})\cap\{x_n=0\}$.*
- (ii) *There is an oriented hyperplane $\Pi\subset\mathbb{R}^n$ parallel to $\vec{\ell}_n$ such that: it meets all the unbounded edges of $\mathcal{K}$ which are nonparallel to $\vec{\ell}_n$, all the vertices of $\mathcal{K}$ and all the unbounded edges of $\mathcal{K}$ parallel to $\vec{\ell}_n$ are contained in $\operatorname{Int}\Pi^-$ and $\pi_n(\mathcal{K}\cap\Pi)=(\mathcal{K}\cap\Pi)\cap\{x_n=0\}$.*

Proof. Denote by $p_1,\dots,p_r$ the vertices of $\mathcal{K}$ and by $\mathcal{A}_1,\dots,\mathcal{A}_s$ the unbounded edges of $\mathcal{K}$, ordered in such a way that the first k (where $0\leq k\leq s$) are nonparallel to $\vec{\ell}_n$ and the remaining $s-k$ are parallel to $\vec{\ell}_n$; for each $i=1,\dots,s$ we write $\mathcal{A}_i:=q_i\vec{v}_i$ where q_i is a vertex of $\mathcal{K}$ and $\vec{v}_i\in\vec{\mathfrak{C}}(\mathcal{K})$ for $i=1,\dots,s$.

Assume first that Λ is bounded. We begin proving (i), that is, $\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))=\vec{\mathfrak{C}}(\mathcal{K})\cap\{x_n=0\}$. Write $H:=\{x_n=0\}$ and let us see that if $\vec{v}:=(v_1,\dots,v_n)\in\vec{\mathfrak{C}}(\mathcal{K})$, then

$$\vec{\pi}_n(\vec{v})=\vec{w}:=(v_1,\dots,v_{n-1},0)\in\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H}.$$

Indeed, if $\vec{v}\in\vec{\ell}_n$, then $\vec{\pi}_n(\vec{v})=\mathbf{0}\in\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H}$; hence, assume that $\vec{v}\not\in\vec{\ell}_n$. Let $p\in\mathcal{K}$; since $\vec{v}\in\vec{\mathfrak{C}}(\mathcal{K})$, then the half-line $p\vec{v}\subset\mathcal{K}$. Since $\vec{v}\not\in\vec{\ell}_n$, then $\vec{w}\in\vec{H}\setminus\{\mathbf{0}\}$. Since the set Λ is bounded and $\pi_n(p\vec{v})=\pi_n(p)\vec{w}$ is a half-line, there is a point $q\in p\vec{v}$ such that the half-line $\pi_n(q\vec{v})=\pi_n(q)\vec{w}$ does not meet $\Lambda\times\{0\}\subset H$. Since for each $t>0$ the point $q+t\vec{v}\in\mathcal{K}\cap\pi_n^{-1}(\pi_n(q)+t\vec{w})$ and $\pi_n(q)+t\vec{w}\not\in\Lambda\times\{0\}$, we deduce that the half-line $\pi_n(q)\vec{w}\subset\mathcal{K}$; hence $\vec{w}\in\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H}$. Thus,

$$\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H}=\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H})\subset\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))\subset\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H},$$

that is, $\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))=\vec{\mathfrak{C}}(\mathcal{K})\cap\{x_n=0\}$. To show (ii) we distinguish two cases:

CASE 1. If $k=0$, all the edges $\mathcal{A}_1,\dots,\mathcal{A}_s$ are parallel to the line $\vec{\ell}_n$. Then $\mathcal{K}$ is in FU-position with respect to the (vectorial) hyperplane $\{x_n=0\}$ suitably oriented and by Remark 2.2 the projection $\pi_n(\mathcal{K})$ is a bounded convex polyhedron. Thus, we can choose a hyperplane Π parallel to the line $\vec{\ell}_n$ such that $\mathcal{K}\subset\operatorname{Int}\Pi^-$; now it is straightforward to show that Π enjoys all the required conditions.

CASE 2. If $k>0$, consider the cone $\mathfrak{C}:=\{\mathbf{0}\}+\vec{\mathfrak{C}}(\mathcal{K})$ which by Lemma 1.7 is a nondegenerated convex polyhedron whose unique vertex is $\mathbf{0}$. We claim that $\pi_n(\mathfrak{C})=\mathfrak{C}\cap\{x_n=0\}$ is also a nondegenerated convex polyhedron whose unique vertex is the origin.

Indeed, by Corollary 1.8, $\vec{\mathfrak{C}}(\mathcal{K})=\{\sum_{i=1}^s\lambda_i\vec{v}_i:\ \lambda_i\geq0\}$. Thus, if we denote $\vec{w}_i=\vec{\pi}_n(\vec{v}_i)$, we have

$$\pi_n(\mathfrak{C})=\mathbf{0}+\Big\{\sum_{i=1}^s\lambda_i\vec{w}_i:\ \lambda_i\geq0\Big\}=\mathbf{0}+\Big\{\sum_{i=1}^k\lambda_i\vec{w}_i:\ \lambda_i\geq0\Big\},$$

which is a convex polyhedron that it is either degenerate or its unique vertex is the origin. Let us see that $\mathbf{0}$ is a vertex of $\pi_n(\mathfrak{C})$. Indeed, let E be a supporting hyperplane of $\mathfrak{C}$

such that $E \cap \mathfrak{C} = \{\mathbf{0}\}$. Since $\pi_n(\mathfrak{C}) \neq \{\mathbf{0}\}$ by (i), we deduce that $E \neq \{x_n = 0\}$ and $E \cap \{x_n = 0\} \neq \varnothing$. Thus, $W := E \cap \{x_n = 0\}$ is a hyperplane of $\{x_n = 0\}$ that meets $\pi_n(\mathfrak{C}) = \mathfrak{C} \cap \{x_n = 0\}$ just in the point $\mathbf{0}$; hence, $\mathbf{0}$ is a vertex of $\mathfrak{C} \cap \{x_n = 0\} = \pi_n(\mathfrak{C})$.

Next, let $\Pi_0 := \mathbf{0} + (\vec{W} + \vec{\ell}_n)$ and let $h_0 \in \mathbb{R}[\mathtt{x}]$ be a degree one polynomial such that $\Pi_0 = \{h_0 = 0\}$; observe that Π_0 is a supporting hyperplane of $\pi_n(\mathfrak{C})$ such that $\Pi_0 \cap \pi_n(\mathfrak{C}) = \{\mathbf{0}\}$. Moreover, $\vec{w}_i \neq 0$ for each $i = 1, \dots, k$, because the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$ of $\mathcal{K}$ are nonparallel to the line $\vec{\ell}_n$. Since Π_0 is a supporting hyperplane for $\pi_n(\mathfrak{C})$ and $\Pi_0 \cap \pi_n(\mathfrak{C}) = \{\mathbf{0}\}$, we may assume that $\vec{h}_0(\vec{w}_i) > 0$ for $i = 1, \dots, k$. Write $\vec{v}_i = \vec{w}_i + t_i \vec{e}_n$ for some $t_i \in \mathbb{R}$; since $\vec{h}_0(\vec{e}_n) = 0$, we deduce that

$$\vec{h}_0(\vec{v}_i) = \vec{h}_0(\vec{w}_i + t_i \vec{e}_n) = \vec{h}_0(\vec{w}_i) + t_i \vec{h}_0(\vec{e}_n) = \vec{h}_0(\vec{w}_i) > 0$$

Let $M > 0$ be such that the ball $\mathcal{B}_n(\mathbf{0}, M)$ contains all the bounded faces of $\mathcal{K}$ and the set $\Lambda \times \{0\}$. Let Π be a hyperplane parallel to Π_0 of equation $h = \vec{h_0} + \rho$ such that $\mathcal{B}_n(\mathbf{0}, M) \subset \{h < 0\}$; in particular, the vertices p_i which are the boundaries of the unbounded edges of $\mathcal{K}$, satisfy $h(p_i) < 0$. Nevertheless, $\vec{h}(\vec{v}_i) = \vec{h}_0(\vec{v}_i) > 0$, and so Π meets the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$. Note that since $\vec{\ell}_n \subset \vec{\Pi}$, then

$$\pi_n^{-1}(\Lambda \times \{0\}) \subset \pi_n^{-1}(\pi_n(\mathcal{B}_n(\mathbf{0}, M))) \subset \pi_n^{-1}(\pi_n(\{h < 0\})) = \{h < 0\}$$

and so, $\pi_n(x) \in \mathcal{K} \cap \Pi \cap \{x_n = 0\}$, for each $x \in \mathcal{K} \cap \{h = 0\} = \mathcal{K} \cap \Pi$ because $\mathcal{K} \cap \Pi \cap \pi_n^{-1}(\Lambda \times \{0\}) = \varnothing$. Thus, we conclude that $\pi_n(\mathcal{K} \cap \Pi) = (\mathcal{K} \cap \Pi) \cap \{x_n = 0\}$.

Assume now that statements (i) and (ii) hold. We distinguish two cases to prove that Λ is bounded:

CASE 1. If $\mathcal{K} \cap \Pi = \varnothing$, o equivalently, all the unbounded edges of $\mathcal{K}$ are parallel to the line $\vec{\ell}_n$. Then $\mathcal{K}$ is in FU-position with respect to the (vectorial) hyperplane $\{x_n = 0\}$ suitably oriented and by Remark 2.2 the projection $\pi_n(\mathcal{K})$ is a bounded convex polyhedron. Thus, $\Lambda \times \{0\} \subset \pi_n(\mathcal{K})$ is a bounded set.

CASE 2. If $\mathcal{K} \cap \Pi \neq \varnothing$, by Lemma 2.1, we have $\mathcal{K} \cap \Pi^+ = (\mathcal{K} \cap \Pi) + \vec{\mathfrak{C}}(\mathcal{K})$,

$$\vec{\mathfrak{C}}(\mathcal{K} \cap \Pi^-) = \Big\{ \sum_{j=k+1}^{s} \lambda_j \vec{v}_j : \ \lambda_j \geq 0 \Big\}$$

and all the vertices of $\mathcal{K}$ and the unbounded edges $\mathcal{A}_{k+1}, \dots, \mathcal{A}_s$ are contained in $\mathcal{K} \cap \operatorname{Int} \Pi^-$. We observe that the vertices of $\mathcal{K} \cap \Pi^-$ correspond to the vertices of $\mathcal{K}$ and the intersections of the edges $\mathcal{A}_1, \dots, \mathcal{A}_k$ with the hyperplane Π. Denote $\{y_j\} = \mathcal{A}_j \cap \Pi$ for $j = 1, \dots, k$. By Corollary 1.8, it holds that $\mathcal{K} \cap \Pi^- = \mathcal{K}_0 + \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi^-)$ where $\mathcal{K}_0$ is the convex hull of the set $\{p_1, \dots, p_r, y_1, \dots, y_k\}$ that it is in particular a bounded convex polyhedron. Since $\vec{\pi}_n(\vec{v}_j) = \mathbf{0}$ for $j = k+1, \dots, s$, we have

$$\pi_n(\mathcal{K} \cap \Pi^-) = \pi_n(\mathcal{K}_0 + \vec{\mathfrak{C}}(\mathcal{K} \cap \Pi^-)) = \pi_n(\mathcal{K}_0) + \vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K} \cap \Pi^-)) = \pi_n(\mathcal{K}_0),$$

which is a bounded set. Thus, if we prove that $\pi_n(\mathcal{K} \cap \Pi^+) = \mathcal{K} \cap \Pi^+ \cap \{x_n = 0\}$, we conclude that $\Lambda \times \{0\} \subset \pi_n(\mathcal{K}_0)$ is bounded. Indeed, by hypothesis it holds

$$\begin{aligned}
\pi_n(\mathcal{K} \cap \Pi^+) = \pi_n((\mathcal{K} \cap \Pi) + \vec{\mathfrak{C}}(\mathcal{K})) &= \pi_n(\mathcal{K} \cap \Pi) + \vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K})) \\
&= (\mathcal{K} \cap \Pi \cap \{x_n = 0\}) + (\vec{\mathfrak{C}}(\mathcal{K}) \cap \{x_n = 0\}) \\
&= ((\mathcal{K} \cap \Pi) + \vec{\mathfrak{C}}(\mathcal{K})) \cap \{x_n = 0\} = \mathcal{K} \cap \Pi^+ \cap \{x_n = 0\},
\end{aligned}$$

and we are done. $\square$

Corollary 2.4 (First trimming position)**.** *Let $\mathcal{K}\subset\mathbb{R}^n$ be a nondegenerated unbounded convex polyhedron and let $\mathcal{F}$ be one of its facets. Then $\mathcal{K}$ is in first trimming position with respect to $\mathcal{F}$ if and only if*

(i) $\mathcal{F}\subset\{x_{n-1}=0\}$ *and* $\mathcal{K}\subset\{x_{n-1}\leq 0\}$,
(ii) $\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))=\vec{\mathfrak{C}}(\mathcal{K})\cap\{x_n=0\}$,
(iii) *There is a hyperplane $\Pi\subset\mathbb{R}^n$ such that: its direction $\vec{\Pi}$ contains the line $\vec{\ell}_n$ generated by $\vec{e}_n$, Π meets all the unbounded edges of $\mathcal{K}$, the vertices of $\mathcal{K}$ are contained in* $\operatorname{Int}\Pi^-$ *and* $\pi_n(\mathcal{K}\cap\Pi)=\mathcal{K}\cap\Pi\cap\{x_n=0\}$.

Proof. We prove first the "only if" implication. The inclusions $\mathcal{F}\subset\{x_{n-1}=0\}$ and $\mathcal{K}\subset\{x_{n-1}\leq 0\}$ are straightforward consequences of the fact that $\mathcal{K}$ is in first trimming position with respecto to the facet $\mathcal{F}$; in particular, $\mathfrak{A}_{\mathcal{K}}:=\{a\in\mathbb{R}^{n-1}:\ \mathcal{I}_a\neq\varnothing,\ (a,0)\notin\mathcal{I}_a\}$ is bounded. By Proposition 2.3, also the two remaining conditions of statement (ii) hold.

Conversely, we have that $\mathfrak{A}_{\mathcal{K}}:=\{a\in\mathbb{R}^{n-1}:\ \mathcal{I}_a\neq\varnothing,\ (a,0)\notin\mathcal{I}_a\}$ is, by Proposition 2.3, bounded, $\mathcal{F}\subset\{x_{n-1}=0\}$ and $\mathcal{K}\subset\{x_{n-1}\leq 0\}$. Moreover, by Remark 2.2 all the intersections $\pi_n^{-1}(a,0)\cap\mathcal{K}\subset((a,0)+\vec{\Pi})\cap\mathcal{K}$ are bounded. Thus, $\mathcal{K}$ is in first trimming position with respect to its facet $\mathcal{F}$. □

Corollary 2.5 (Second trimming position)**.** *Let $\mathcal{K}\subset\mathbb{R}^n$ be a nondegenerated unbounded convex polyhedron and let $\mathcal{F}$ be one of its facets. Then $\mathcal{K}$ is in second trimming position with respect to $\mathcal{F}$ if and only if*

(i) $\mathcal{F}\subset\{x_n=0\}$ *and* $\mathcal{K}\subset\{x_n\leq 0\}$,
(ii) $\vec{\pi}_n(\vec{\mathfrak{C}}(\mathcal{K}))=\vec{\mathfrak{C}}(\mathcal{K})\cap\{x_n=0\}$,
(iii) *There is a hyperplane $\Pi\subset\mathbb{R}^n$ such that: its direction $\vec{\Pi}$ contains the line $\vec{\ell}_n$ generated by $\vec{e}_n$, Π meets all the unbounded edges of $\mathcal{K}$ nonparallel to $\vec{\ell}_n$, the vertices of $\mathcal{K}$ are contained in the open half-space* $\operatorname{Int}\Pi^-$ *and* $\pi_n(\mathcal{K}\cap\Pi)=\mathcal{K}\cap\Pi\cap\{x_n=0\}$.

Proof. We begin with the "only if" implication. It is clear that $\mathcal{F}\subset\{x_n=0\}$ and $\mathcal{K}\subset\{x_n\leq 0\}$; moreover $\mathfrak{A}_{\mathcal{K}}:=\{a\in\mathbb{R}^{n-1}:\ \mathcal{I}_a\neq\varnothing,\ (a,0)\notin\mathcal{I}_a\}$ is bounded. By Proposition 2.3, the remaning conditions hold.

Conversely, we have that $\mathfrak{A}_{\mathcal{K}}:=\{a\in\mathbb{R}^{n-1}:\ \mathcal{I}_a\neq\varnothing,\ (a,0)\notin\mathcal{I}_a\}$ is bounded, $\mathcal{F}\subset\{x_n\leq 0\}$ and $\mathcal{K}\subset\{x_n\leq 0\}$. Besides, since $\mathcal{K}\subset\{x_n=0\}$, all the intersections $\pi_n^{-1}(a,0)\cap\mathcal{K}\subset((a,0)+\vec{\Pi})\cap\mathcal{K}$ are upperly bounded. Therefore, $\mathcal{K}$ is in second trimming position with respect to its facet $\mathcal{F}$. □

3. PLACING IN TRIMMING POSITIONS

3.1. **First trimming position.** The purpose of the first part of this section is to prove Theorem 1. For the sake of clearness, we divide the proof of Theorem 1 into two parts.

Proof of Theorem 1 (i). By Lemma 1.4, we may assume that $\mathcal{K}$ is in FU-position with respect to the hyperplane $\vec{\Pi}=\{x_n=0\}$. The proof runs in several steps (see Figure 3):

Step 1. We begin by choosing the facet $\mathcal{F}$ of the statement. Denote $\widehat{\mathcal{K}}:=\operatorname{Cl}_{\mathbb{RP}^3}(\mathcal{K})$ and $\mathcal{K}_\infty:=\operatorname{Cl}_{\mathbb{RP}^3}(\mathcal{K})\cap\mathsf{H}_\infty(\mathbb{R})$; clearly, $\widehat{\mathcal{K}}=\mathcal{K}\sqcup\mathcal{K}_\infty$. Consider the involutive homography

$$\phi:\mathbb{RP}^3\to\mathbb{RP}^3,\ (x_0:x_1:x_2:x_3)\mapsto(x_3:x_1:x_2:x_0)=(y_0:y_1:y_2:y_3).$$

By Lemma 1.6, the convex polyhedron $\mathcal{K}':=\phi(\widehat{\mathcal{K}})\subset\mathbb{R}^3\equiv\{y_0\neq 0\}$ is bounded and one of its faces is $\mathcal{E}':=\phi(\mathcal{K}_\infty)$, which needs not to be a facet. Now we distinguish:

Case 1. If $\mathcal{E}'$ has dimension 0 or 1, we choose an unbounded facet $\mathcal{F}$ of $\mathcal{K}$ such that $\mathcal{E}' \subset \operatorname{Cl}(\phi(\mathcal{F}))$.

Case 2. If $\mathcal{E}'$ has dimension 2 (that is, $\mathcal{E}'$ is a facet of $\phi(\widehat{\mathcal{K}})$), we choose an unbounded facet $\mathcal{F}$ of $\mathcal{K}$ such that $\dim(\operatorname{Cl}(\phi(\mathcal{F})) \cap \mathcal{E}') = 1$.

Step 2. Denote by H the plane generated by the facet $\mathcal{F}$ and let $\mathfrak{F}$ be the (finite) set constituted by the intersections of H with the unbounded edges of $\mathcal{K}$ which are nonparallel to H. Since $\mathcal{K}$ is in FU-position with respect to $\vec{\Pi}$ we may choose an oriented plane Π parallel to $\vec{\Pi}$ that meets all the unbounded edges and such that all the vertices of $\mathcal{K}$ are contained in the half-space Π^-. Moreover, if $\dim(\mathcal{E}') = 2$, then the dimension of $\mathcal{K}_\infty$ is 2, that of $\operatorname{Cl}_{\mathbb{RP}^3}(\mathcal{F}) \cap \mathcal{K}_\infty$ is 1 and, by Lemma 1.7, the dimension of $\vec{\mathfrak{C}}(\mathcal{F})$ is 2. Since $\mathcal{K}$ is in FU-position with respect to $\vec{\Pi}$, then $\mathcal{F}$ is in FU-position with respect to $\vec{\Pi} \cap \vec{H}$ inside H. Fix $\vec{w} \in \operatorname{Int}(\vec{\mathfrak{C}}(\mathcal{F}))$ and, by Lemma 1.9 (ii), we may assume that the intersection $p\vec{w} \cap (\mathcal{F} \cap \Pi) = p\vec{w} \cap (H \cap \mathcal{K} \cap \Pi)$ is a singleton for each $p \in \mathfrak{F}$.

Step 3. Denote $\mathcal{P} := \mathcal{K} \cap \Pi$ and $\mathcal{E}$ the intersection $\mathcal{F} \cap \Pi = H \cap \mathcal{P}$. Consider the quotient map $\rho : \Pi \to \Pi/(\vec{H} \cap \vec{\Pi})$, $x \mapsto x + (\vec{H} \cap \vec{\Pi})$ which is continuous with respect to the quotient topology of $\Pi/(\vec{H} \cap \vec{\Pi})$. Since $\mathcal{P} \subset \Pi$ is compact and $\Pi/(\vec{H} \cap \vec{\Pi})$ is homeomorphic to $\mathbb{R}$ with the usual topology, $\rho(\mathcal{P}) \equiv [a, b]$ is a closed bounded interval (nontrivial because $\mathcal{P}$ has dimension 2). Also $\Lambda_a := H \cap \Pi$ is a supporting line of $\mathcal{P}$ in Π, because it contains one of its edges, and we may assume that $\rho^{-1}(a) \cap \mathcal{P} = H \cap \mathcal{P} \subset \Lambda_a$. Observe that $\Lambda_b := \rho^{-1}(b)$, which is a line contained in Π parallel to $H \cap \Pi$, is a supporting line of $\mathcal{P}$ in Π; hence, $\Lambda_b \cap \mathcal{P}$ is a face of $\mathcal{P}$, that is, an edge or a vertex. Since $\mathcal{P}$ is a bounded polygon, there is a vertex $p_0 \in \Lambda_b \cap \mathcal{P}$; by Remark 2.2 we know that $\{p_0\}$ is the intersection of Π with an unbounded edge $\mathcal{A}$ of $\mathcal{K}$.

Let H_0 be the plane of $\mathbb{R}^3$ that contains Λ_b and the edge $\mathcal{A}$. By Lemma 1.5, we have that H_0 is a supporting plane of $\mathcal{K}$.

Step 4. Let ℓ be the line generated by $\mathcal{A}$; we are going to construct a plane W that contains ℓ and such that $W \cap H \cap \mathcal{K} = W \cap \mathcal{F}$ is a half-line. We analyze two possible situations:

Case 1. If the line ℓ is parallel to the plane H, we choose a plane W of $\mathbb{R}^3$ that contains ℓ and meets $\operatorname{Int} \mathcal{F} \cap \Pi \subset H \cap \mathcal{P}$ in a point q.

Case 2. If the line ℓ is nonparallel to H, then it meets H in a singleton, say $\ell \cap H = \{q_0\}$; besides,

$$\operatorname{Cl}_{\mathbb{RP}^3}(\ell) \cap \operatorname{Cl}_{\mathbb{RP}^3}(H) \cap \mathsf{H}_\infty(\mathbb{R}) = \varnothing \implies \operatorname{Cl}_{\mathbb{RP}^3}(\mathcal{A}) \cap \operatorname{Cl}_{\mathbb{RP}^3}(\mathcal{F}) \cap \mathcal{K}_\infty = \varnothing$$

and so, the dimension of $\mathcal{K}_\infty$ is 2. Recall now the vector $\vec{w}$ fixed in Step 2; by our choice of $\mathcal{F}$ and Π (see Steps 1 and 2) the intersection

$$q_0\vec{w} \cap (\mathcal{F} \cap \Pi) = q_0\vec{w} \cap (H \cap \mathcal{K} \cap \Pi)$$

is a point q. The line through q_0 that is parallel to $\vec{w}$ is coplanar to ℓ; denote by W the plane of $\mathbb{R}^3$ that contains both lines.

Step 5. Let $\pi : \mathbb{R}^3 \to \mathbb{R}^3$ be the projection onto W in the direction of $\vec{H} \cap \vec{\Pi}$. Observe that in both cases, since $p_0 \in \Lambda_b \cap \Pi$, $q \in \Lambda_a \cap \Pi$, and these lines are parallel to the line $\vec{H} \cap \vec{\Pi}$, the line $W \cap \Pi$ (through the points p_0 and q) satisfies $\pi(\mathcal{K} \cap \Pi) = (\mathcal{K} \cap \Pi) \cap W$. Moreover, we also have that $\vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K})) = \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W}$.



(a) (b)

FIGURE 3. First trimming position for unbounded convex polyhedra.

Indeed, let us write $H_0 := \{h_0 = 0\}$, $H := \{h = 0\}$, where $h, h_0 \in \mathbb{R}[\mathtt{x}]$ are degree one polynomials, and $\vec{H}_0^+ := \{\vec{h}_0 \geq 0\}$, $\vec{H}^+ := \{\vec{h} \geq 0\}$. Since $\mathcal{K} \subset H_0^+ \cap H^+$, we have

$$\vec{\mathfrak{C}}(\mathcal{K}) \subset \vec{\mathfrak{C}}(H_0^+ \cap H^+) = \vec{H}_0^+ \cap \vec{H}^+.$$

Next, we set $\mathcal{A} := q_1\vec{w}_1$ and $W \cap H \cap \mathcal{K} := q_2\vec{w}_2$, where $q_1, q_2 \in \mathcal{K}$ and $\vec{w}_1, \vec{w}_2 \in \vec{\mathfrak{C}}(\mathcal{K}) \setminus \{\mathbf{0}\}$. We distinguish now two cases:

CASE 1. If $\vec{w}_1$ and $\vec{w}_2$ are linearly dependent, then H_0 and H are planes parallel to $\vec{H}_0^+ \cap \vec{H}^+ = \vec{H}$. Therefore,

$$\{\lambda\vec{w}_1 : \ \lambda \geq 0\} \subset \vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K})) \subset \vec{H} \cap \vec{W} = \{\mu\vec{w}_1 : \ \mu \in \mathbb{R}\},$$

and since $\mathcal{K}$ is nondegenerated $\vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K})) = \{\lambda\vec{w}_1 : \ \lambda \geq 0\} = \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W}$.

CASE 2. If $\vec{w}_1$ and $\vec{w}_2$ are linearly independent,

$$\vec{H}_0^+ \cap \vec{H}^+ \cap \vec{W} = \{\lambda_1\vec{w}_1 + \lambda_2\vec{w}_2 : \ \lambda_1, \lambda_2 \geq 0\} = \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W}.$$

Thus,

$$\vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W} = \vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W}) \subset \vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K})) \subset \vec{H}_0^+ \cap \vec{H}^+ \cap \vec{W} = \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W},$$

that is, $\vec{\pi}(\vec{\mathfrak{C}}(\mathcal{K})) = \vec{\mathfrak{C}}(\mathcal{K}) \cap \vec{W}$.

Step 6. After a change of coordinates that transforms H^+ in $\{x_2 \leq 0\}$, W in $\{x_3 = 0\}$ and the line $\vec{H} \cap \vec{\Pi}$ onto the line $\{x_1 = x_2 = 0\}$, we conclude, by Proposition 2.4, that the convex polyhedron $\mathcal{K}$ is in first trimming position respect to the facet $\mathcal{F}$. $\square$

Next, we prove the second part of Theorem 1.

Proof of Theorem 1 (ii). Let us see first that $\mathcal{K}$ has a bounded facet. We may assume that the unbounded edges of $\mathcal{K}$ are parallel to the vector $\vec{e}_3$. Let $\mathcal{A}$ be an unbounded edge of $\mathcal{K}$ and let $p \in \partial\mathcal{A}$. Let $\mathcal{F}$ be a facet of $\mathcal{K}$ nonparallel to $\vec{e}_3$ such that $p \in \mathcal{F}$. It holds that $\vec{\mathfrak{C}}(\mathcal{F}) \subset \vec{\mathfrak{C}}(\mathcal{K}) = \{\lambda\vec{e}_3 : \ \lambda \geq 0\}$ and since $\mathcal{F}$ is nonparallel to $\vec{e}_3$, we deduce that $\vec{\mathfrak{C}}(\mathcal{F}) = \{\mathbf{0}\}$; hence, $\mathcal{F}$ is bounded.

Fix a bounded facet $\mathcal{F}_0$ of $\mathcal{K}$ and let $H \subset \mathbb{R}^3$ be the plane generated by $\mathcal{F}_0$. Since the unbounded edges of $\mathcal{K}$ are parallel, $\vec{\mathfrak{C}}(\mathcal{K}) = \{\lambda\vec{v} : \ \lambda \geq 0\}$ where $\vec{v} \in \mathbb{R}^3$ is a directional vector of the unbounded edges of $\mathcal{K}$; let $\vec{\ell}$ be the line generated by $\vec{v}$. Since $\mathcal{F}_0$ is a bounded facet and $\mathcal{K} \cap H = \mathcal{F}_0$, then $\vec{H} \cap \vec{\ell} = \{0\}$; hence, a line parallel to $\vec{\ell}$ and a plane parallel to $\vec{H}$ meet in a singleton. Let now Π be a plane parallel to H that meets all the unbounded edges of $\mathcal{K}$ and such that all the bounded faces of $\mathcal{K}$ are contained in $\operatorname{Int}\Pi^-$. We deduce from Remark 2.2 that $\mathcal{P} := \mathcal{K} \cap \Pi$ is a bounded convex polygon, $\mathcal{K} \cap \Pi^-$ is a bounded convex polyhedron and moreover $\mathcal{K} \cap \Pi^+ = \mathcal{P} + \vec{\mathfrak{C}}(\mathcal{K}) = \mathcal{P} + \{\lambda\vec{v} : \ \lambda \geq 0\}$; hence, $\mathcal{K} \cap \Pi^+$ is affinely equivalent to $\mathcal{P} \times [0, +\infty[$. After a change of coordinates, we may assume that:

- H is the plane $\{x_2 = 0\}$ and $\mathcal{K} \subset \{x_2 \leq 0\}$,
- Π is the plane $\{x_2 = -1\}$ and $\pi_3(\mathcal{P}) = \mathcal{P} \cap \{x_3 = 0\}$ (recall here the projection property of convex polygons described in the Introduction),
- $\vec{v} = -\vec{e}_2$ and so $\vec{\mathfrak{C}}(\mathcal{K}) = \{-\lambda\vec{e}_2 : \ \lambda \geq 0\} \subset \{x_3 = 0\}$.

Since $\pi_3(\vec{\mathfrak{C}}(\mathcal{K})) = \vec{\mathfrak{C}}(\mathcal{K}) \cap \{x_3 = 0\}$, we deduce from Lemma 2.4 that $\mathcal{K}$ is in first trimming position with respect to the facet $\mathcal{F}_0$. $\square$

3.2. **Second trimming position.** Our purpose now is to prove Theorem 2.

Proof of Theorem 2. (i) After a change of coordinates we may assume that $\mathcal{K} \subset \{x_3 \leq 0\}$ and $\mathcal{K} \cap \{x_3 = 0\} = \mathcal{F}$. Observe that $\mathcal{F}$ is an unbounded convex polygon and so it has exactly two unbounded edges $\mathcal{A}_1$ and $\mathcal{A}_2$ which are nonparallel because $\dim \vec{\mathfrak{C}}(\mathcal{F}) = 2$. Let $\mathcal{F}_1$ and $\mathcal{F}_2$ be the (unbounded) facets of $\mathcal{K}$ that satisfy $\mathcal{A}_i = \mathcal{F}_i \cap \mathcal{F}$. Let H_1 and H_2 be the planes of $\mathbb{R}^3$ generated by $\mathcal{F}_1$ and $\mathcal{F}_2$. Observe that since $H_1 \cap \{x_3 = 0\}$ and $H_2 \cap \{x_3 = 0\}$ are nonparallel lines, we may assume after a change of coordinates that keeps invariant the plane $\{x_3 = 0\}$ that $H_1 := \{x_1 = 0\}$ and $H_2 := \{x_2 = 0\}$. Moreover, after a change of the type $(x_1, x_2, x_3) \mapsto (\pm x_1, \pm x_2, \pm x_3)$ we may assume that $\mathcal{K} \subset \{x_1 \geq 0, x_2 \geq 0, x_3 \leq 0\}$. In particular, the direction of the intersection line $\ell_3 := H_1 \cap H_2$ is the line $\vec{\ell}_3$ generated by the vector $\vec{e}_3$. Consider the projection

$$\pi_3 : \mathbb{R}^3 \to \mathbb{R}^3, (x_1, x_2, x_3) \mapsto (x_1, x_2, 0)$$

onto $H_3 := \{x_3 = 0\}$ in the direction of $\vec{\ell}_3$. Let us see that there is an $M \geq 0$ such that $\mathcal{K}_M := \mathcal{K} \cup]-M, M[^3$ satisfies $\pi_3(\mathcal{K}_M) = \mathcal{K}_M \cap \{x_3 = 0\}$ and then, by the Remarks in the Introduction, we deduce that $\mathcal{K}$ is in second trimming position with respecto to its facet $\mathcal{F}$). Choose $M \geq 0$ such that the vertices p_1 and p_2 of $\mathcal{K}$ which are respectively the boundary points of $\mathcal{A}_1$ and $\mathcal{A}_2$ are contained in $]-M, M[^3$. Note that $\mathcal{A}_1 \subset \{x_1 = 0, x_2 \geq 0, x_3 = 0\}$ and $\mathcal{A}_2 \subset \{x_1 \geq 0, x_2 = 0, x_3 = 0\}$. Since the convex polygon $\mathcal{P}$ of unbounded edges $\mathcal{A}_1$ and $\mathcal{A}_2$ and vertices p_1, p_2 is contained in $\mathcal{F}$, we deduce that

$$\mathcal{F} \cup (]-M, M[^2 \times \{0\}) = \{x_1 \geq 0, x_2 \geq 0, x_3 = 0\} \cup (]-M, M[^2 \times \{0\}).$$

Thus, since $\mathcal{K}\subset\{x_1\geq 0,x_2\geq 0,x_3\leq 0\}$ we have

$$\begin{aligned}\{x_1\geq 0,x_2\geq 0,x_3=0\}\cup(\,]-M,M[^2\times\{0\})&\subset\pi_3(\mathcal{K}\cup\,]-M,M[^3)\\ &\subset\pi_3(\{x_1\geq 0,x_2\geq 0,x_3\leq 0\})\cup(\,]-M,M[^2\times\{0\})\\ &=\{x_1\geq 0,x_2\geq 0,x_3=0\}\cup(\,]-M,M[^2\times\{0\})\end{aligned}$$

and in this way

$$\begin{aligned}\pi_3(\mathcal{K}_M)&=\{x_1\geq 0,x_2\geq 0,x_3=0\}\cup(\,]-M,M[^2\times\{0\})\\ &=\mathcal{F}\cup(\,]-M,M[^2\times\{0\})=\mathcal{K}_M\cap\{x_3=0\},\end{aligned}$$

as wanted.

(ii) First, observe that if $\mathcal{F}$ is a bounded facet of $\mathcal{K}$, then $\mathcal{F}$ is nonparallel to the unbounded edges of $\mathcal{K}$. Otherwise, let H be the plane generated by $\mathcal{F}$. Then, $\vec{\mathfrak{C}}(\mathcal{K})\subset\vec{H}$ and so

$$\vec{\mathfrak{C}}(\mathcal{K})\cap\vec{H}=\vec{\mathfrak{C}}(\mathcal{K}\cap H)=\vec{\mathfrak{C}}(\mathcal{F})=\{\mathbf{0}\},$$

a contradiction. After a change of coordinates we may assume that $\mathcal{K}\subset\{x_3\leq 0\}$, $\mathcal{K}\cap\{x_3=0\}$ is a face of $\mathcal{K}$ and the unbounded edges of $\mathcal{K}$ are parallel to the vector $\vec{e}_3$. Thus, $\mathcal{K}$ is in FU-position with respect to the hyperplane $\vec{\Pi}:=\{x_3=0\}$ (with orientation opposite to the usual one). By Corollary 1.8 (iii) and using that $\vec{\pi}_3(\vec{e}_3)=\mathbf{0}$, we deduce that $\pi_3(\mathcal{K})$ is a bounded convex polyhedron and so, if we choose $M>0$ such that $\pi_3(\mathcal{K})\subset\,]-M,M[^3\cap\{x_3=0\}$, we deduce that

$$\pi_3(\mathcal{K}\cup\,]-M,M[^3)=\,]-M,M[^3\cap\{x_3=0\}=(\mathcal{K}\cup\,]-M,M[^3)\cap\{x_3=0\}$$

and, as we have seen in the Remarks in the Introduction, we conclude that $\mathcal{K}$ is in second trimming position with respect to its facet $\mathcal{F}$. □

4. Limitations of the trimming positions

The purpose of this section is to provide examples of unbounded convex polyhedra which present difficulties in order to be placed in first or in second trimming position with respect to their facets. We start by defining a sectional projection:

Definition 4.1. Let $\mathcal{K}$ be a convex polyhedron in $\mathbb{R}^n$, and $H\subset\mathbb{R}^n$ a hyperplane. We say that $\mathcal{K}\cap H$ is a *sectional projection of* $\mathcal{K}$ if there exists a linear projection $\pi:\mathbb{R}^n\to H$ with direction given by a line $\vec{\ell}$ (nonparallel to H) such that $\pi(\mathcal{K})=\mathcal{K}\cap H$.

Our key result is the following:

Theorem 4.2 (Convex polyhedra without sectional projections). *For each $n\geq 3$ there is an n-dimensional bounded convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ wich does not have sectional projections.*

Corollary 4.3 (Counterexamples to the trimming positions). *Let $n\geq 4$ and $d:=n-1$ and let $\mathcal{P}\subset\mathbb{R}^d$ be a bounded convex polyhedron without sectional projections. Define $\mathcal{K}:=\mathcal{P}\times[0,+\infty[\,\subset\mathbb{R}^n$. Then $\mathcal{K}$ cannot be placed in first trimming position with respect to any of its facets. Moreover, $\mathcal{K}$ cannot be placed in second trimming position with respect to any of its unbounded facets.*

Proof. Denote $\mathcal{F}:=\mathcal{K}\cap\{x_n=0\}=\mathcal{P}\times\{0\}$ which is a facet of $\mathcal{K}$ and $\vec{\ell}$ the direction of the unbounded edges of $\mathcal{K}$. Suppose first that, after a change of coordinates, $\mathcal{K}$ is placed in first trimming position with respect to one of its facets. By Corollary 2.4 there is a hyperplane $\Pi \subset \mathbb{R}^n$ such that: its direction $\vec{\Pi}$ contains the line $\vec{\ell_n}$ generated by $\vec{e}_n$, Π meets each unbounded edge of $\mathcal{K}$ in a singleton, the vertices of $\mathcal{K}$ are contained in $\operatorname{Int}\Pi^-$ and $\pi_n(\mathcal{K}\cap\Pi) = \mathcal{K}\cap\Pi\cap\{x_n=0\}$.

(4.3.1) Consider the projection $\rho : \mathbb{R}^n \to \mathbb{R}^n$ in the direction of the line $\vec{\ell}$ onto the hyperplane H generated by $\mathcal{F}$. Since Π meets the unbounded edges of $\mathcal{K}$ in a singleton and all of them are parallel to the line $\vec{\ell}$, we have that Π is nonparallel to $\vec{\ell}$; hence, $g := \rho|_{\Pi} : \Pi \to H$ is an affine bijection. Moreover, since $\vec{\rho}(\vec{\ell}) = \{\mathbf{0}\}$ and by Remark 2.2, $\mathcal{P}_1 := \mathcal{K}\cap\Pi$ is the bounded convex polyhedron whose vertices are the intersections of the unbounded edges of $\mathcal{K}$ (all of them parallel to the line $\vec{\ell}$) with the hyperplane Π, we deduce that $\rho(\mathcal{P}_1)$ is the bounded convex polyhedron whose vertices are the intersections of the unbounded edges of $\mathcal{K}$ with the plane H, that is, $\rho(\mathcal{P}_1) = \mathcal{F}$. Now, if we define $\vec{r} := \vec{g}(\vec{\ell_n})$ and $W := g(\Pi\cap\{x_n=0\})$, then we have that the projection $\pi := (g\circ\pi_n\circ g^{-1}) : H \to H$ in the direction of $\vec{r}$ onto the hyperplane W satisfies $\pi(\mathcal{F}) = \mathcal{F}\cap W$, but this contradicts the fact that $\mathcal{P}$ has no sectional projections because $\mathcal{P}$ and $\mathcal{F}$ are affinely equivalent. Thus, $\mathcal{K}$ cannot be placed in first trimming position with respect to any of its facets.

Suppose next that, after a change of coordinates, $\mathcal{K}$ can be placed in second trimming position with respect to one of its unbounded facets $\mathcal{F}$, that now is contained in the hyperplane $\{x_n = 0\}$. In particular, the unbounded edges of $\mathcal{K}$ are parallel to a line $\vec{\ell}$ contained in the hyperplane $\{x_n = 0\}$, and so they are nonparallel to the line $\vec{\ell_n}$ generated by $\vec{e}_n$. By Corollary 2.5, there is a hyperplane $\Pi \subset \mathbb{R}^n$ such that: its direction $\vec{\Pi}$ contains the line $\vec{\ell_n}$, the hyperplane Π meets all the unbounded edges of $\mathcal{K}$ nonparallel to $\vec{\ell_n}$ (that is, all the unbounded edges of $\mathcal{K}$, because none of them is parallel to $\vec{\ell_n}$), the bounded edges of $\mathcal{K}$ are contained in the open half-space $\operatorname{Int}\Pi^-$ and $\pi_n(\mathcal{K}\cap\Pi) = \mathcal{K}\cap\Pi\cap\{x_n=0\}$. Proceeding analogously to 4.3.1, we arrive at a contradiction. Thus, $\mathcal{K}$ cannot be placed in second trimming position with respect to any of its unbounded facets. □

4.1. **Strategy.** The following result will help us to show Theorem 4.2.

Lemma 4.4. *Let $\mathcal{K} \subset \mathbb{R}^n$ be a convex polyhedron of dimension $n \geq 3$ and let $\pi : \mathbb{R}^n \to \mathbb{R}^n$ be the projection in the direction of the line $\vec{\ell}$ onto the hyperplane H. Let $p_1, \dots, p_\ell$ be the vertices of $\mathcal{K}$ contained in H and let $\mathcal{A}_1, \dots, \mathcal{A}_k$ be the edges of $\mathcal{K}$ parallel to $\vec{\ell}$. Suppose that $\mathcal{P} := \pi(\mathcal{K}) = \mathcal{K}\cap H$ and write $\{q_j\} = \pi(\mathcal{A}_j)$ for $j = 1, \dots, k$. Then the vertices of $\mathcal{P}$ are $q_1, \dots, q_k$ and $p_1, \dots, p_\ell$. In particular,*

- (i) *If $\vec{\ell}$ is parallel to none of the edges of $\mathcal{K}$, then the vertices of $\mathcal{P}$ are $p_1, \dots, p_\ell$.*
- (ii) *If $\mathcal{K}$ has pairwise nonparallel edges, then $\mathcal{P}$ has at most $\ell + 1$ vertices.*

The proof of the previous lemma requires some preparation:

Lemma 4.5. *Let $\mathcal{K} \subset \mathbb{R}^n$ be a convex polyhedron of dimension $n \geq 3$, let $\mathcal{A}$ be an edge of $\mathcal{K}$, let ℓ be the line generated by $\mathcal{A}$ and let $\pi : \mathbb{R}^n \to \mathbb{R}^n$ be the projection in the direction of the line $\vec{\ell}$ onto the hyperplane $H \subset \mathbb{R}^n$. Let E be a supporting hyperplane of $\mathcal{K}$ such that $E\cap\mathcal{K} = \mathcal{A}$. Then $E\cap H$ is a supporting hyperplane of $\mathcal{P} := \pi(\mathcal{K})$ in H such that $E\cap\mathcal{P} = H\cap\ell$ is a vertex of $\mathcal{P}$.*

Proof. First, since $\dim\mathcal{K} = n$, we deduce that $\mathcal{P} := \pi(\mathcal{K})$ has dimension $n-1 \geq 2$; moreover, since $\vec{\ell} \not\subset \vec{H}$, then $H\cap\ell = \{p\}$. Thus, if we prove that $E\cap\mathcal{P} = H\cap\ell$, we deduce that $E\cap H$ is a supporting hyperplane of $\mathcal{P} := \pi(\mathcal{K})$ in H and that p is a vertex of $\mathcal{P}$. Indeed, since $\ell \subset E$, it holds $\pi(E) = E\cap H$ and $E = \pi^{-1}(\pi(E))$; hence, if $y \in E\cap\mathcal{P} = \pi(E)\cap\mathcal{P}$

there is $x \in \mathcal{K}$ such that $\pi(x) = y$ and so $x \in \pi^{-1}(y) \subset \pi^{-1}(\pi(E)) = E$. This means that $x \in E \cap \mathcal{K} = \mathcal{A}$ and therefore $y = \pi(x) \in \pi(\mathcal{A}) = H \cap \ell = \{p\}$. □

Lemma 4.6. *Let $\mathcal{K} \subset \mathbb{R}^n$ be a convex polyhedron of dimension $n \geq 3$ and let $\pi : \mathbb{R}^n \to \mathbb{R}^n$ be the projection in the direction of the line $\vec{\ell}$ onto the hyperplane H. Let $p_1, \dots, p_r$ be the vertices of $\mathcal{K}$ (if any) and let $\mathcal{A}_1, \dots, \mathcal{A}_k$ be the edges of $\mathcal{K}$ parallel to $\vec{\ell}$ (if any). Then*

- (i) *$\pi(\mathcal{A}_1), \dots, \pi(\mathcal{A}_k)$ are vertices of $\mathcal{P} := \pi(\mathcal{K})$.*
- (ii) *If p is a vertex of $\mathcal{P}$ such that $p \notin \pi(\mathcal{A}_i)$ for $i = 1, \dots, k$, and W is a supporting hyperplane of $\mathcal{P}$ in H with $\mathcal{P} \cap W = \{p\}$, then there is $j = 1, \dots, r$ such that $\pi(p_j) = p$. Moreover, the hyperplane $\Pi := p_j + \vec{W} + \vec{\ell}$ is a supporting hyperplane of $\mathcal{K}$ such that $\Pi \cap \mathcal{K} = \{p_j\}$.*

Proof. First, by Lemma 4.5, $\pi(\mathcal{A}_1), \dots, \pi(\mathcal{A}_k)$ provide vertices of $\mathcal{P} := \pi(\mathcal{K})$, and so (i) is proved.

(ii) Let now p be one of the remaining vertices of $\mathcal{P}$ (if any) and let W be a supporting hyperplane of $\mathcal{P}$ in H such that $W \cap \mathcal{P} = \{p\}$. After a change of coordinates, we may assume that $H := \{x_n = 0\}$, $\vec{\ell}$ is the line generated by the vector $\vec{e}_n$, $p := \mathbf{0}$, $W := \{x_{n-1} = 0\} \cap H$ and $\mathcal{P} \subset \{x_{n-1} \geq 0\} \cap H$.

Let Π be the hyperplane generated by W and the line $p + \vec{\ell}$, that is, $\Pi := \{x_{n-1} = 0\}$. Observe that

$$\mathcal{K} \subset \pi^{-1}(\mathcal{P}) \subset \pi^{-1}(\{x_{n-1} \geq 0\} \cap H) = \{x_{n-1} \geq 0\} = \Pi^+.$$

To show that Π is a supporting hyperplane of $\mathcal{K}$ it is enough to check that Π meets $\mathcal{K}$. To that end, observe that if $x \in \mathcal{K}$ satisfies $\pi(x) = p \in W$, then $x \in \pi^{-1}(W) = \Pi$.

Now, we have that $\mathcal{E} := \mathcal{K} \cap \Pi$ is a face of $\mathcal{K}$ satisfying $p \in \pi(\mathcal{E}) = \pi(\mathcal{K} \cap \Pi) \subset \mathcal{P} \cap W = \{p\}$; hence, $\pi(\mathcal{E}) = \{p\}$. Thus, $\mathcal{E}$ either has dimension 0 or has dimension 1, being in this last case parallel to $\vec{\ell}$. But since $\{p\}$ is not the projection of an edge of $\mathcal{K}$ parallel to $\vec{\ell}$, we deduce that $\dim \mathcal{E} = 0$ and so it is a vertex p_j of $\mathcal{K}$. This means that $p = \pi(p_j)$ and $\mathcal{K} \cap \Pi = \{p_j\}$, as wanted. □

Proof of Lemma 4.4. First, by Lemma 4.6, $q_1, \dots, q_k$ are vertices of $\mathcal{K}$. On the other hand, observe that $\mathcal{P}$ is a convex polyhedron of dimension $n - 1 \geq 2$. Let Π_i be a supporting hyperplane of $\mathcal{K}$ such that $\Pi_i \cap \mathcal{K} = \{p_i\}$; since $\dim \mathcal{P} \geq 2$ and $p_i \in H$, we deduce that $\Pi_i \neq H$ is nonparallel to H. Define $W_i := \Pi_i \cap H$, which is a hyperplane of H; since $p_i \in H$, then $W_i \cap \mathcal{P} = (\Pi_i \cap H) \cap (\mathcal{K} \cap H) = \{p_i\}$ and so p_i is a vertex of $\mathcal{P}$.

Let now p be a vertex of $\mathcal{P}$ different from $q_1, \dots, q_k$. Choose a hyperplane W of H such that $W \cap \mathcal{P} = \{p\}$. By Lemma 4.6, there is a vertex q of $\mathcal{K}$ such that $\pi(q) = p$; moreover, the hyperplane $\Pi := q + \vec{W} + \vec{\ell}$ is a supporting hyperplane of $\mathcal{K}$ such that $\Pi \cap \mathcal{K} = \{q\}$. Since $\mathcal{P} = \mathcal{K} \cap H$, then $p \in \mathcal{K}$ and so the segment $\overline{pq} \subset \mathcal{K}$; moreover, such segment is contained in the line $q + \vec{\ell} \subset \Pi$. Thus, $\overline{pq} \subset \mathcal{K} \cap \Pi = \{q\}$, that is, $p = q$ is a vertex of $\mathcal{K}$ contanided in H and so $p = p_i$ for some $i = 1, \dots, \ell$.

Once this is proved, statements (i) and (ii) are straightforward. □

4.2. **Spherical polyhedra.** An n-dimensional convex polyhedron $\mathcal{K} \subset \mathbb{R}^n$ is called *spherical* if it is the convex hull of a finite subset $\mathfrak{F} := \{p_1, \dots, p_r\}$ $(r \geq n + 1)$ of the unitary sphere $\mathbb{S}^{n-1} := \{x_1^2 + \cdots + x_n^2 = 1\}$ centered at the origin. Since the tangent hyperplane to $\mathbb{S}^{n-1}$ at the point p_i meets $\mathcal{K}$ just in the point p_i, the points p_i are vertices of $\mathcal{K}$. Conversely, the set of vertices of the convex hull $\mathcal{K}$ of the finite set $\mathfrak{F}$ is contained in $\mathfrak{F}$; hence, the set of vertices of $\mathcal{K}$ coincides with $\mathfrak{F}$. Moreover, since the sphere $\mathbb{S}^{n-1}$ is convex,

$\mathcal{K}\subset\mathbb{S}^{n-1}$ and since $\mathcal{K}$ is n-dimensional, the set $\mathfrak{F}$ contains $n+1$ points that are affinely independent. The spherical polyhedron $\mathcal{K}$ is *generic* if each subset of $n+1$ vertices of $\mathcal{K}$ is affinely independent. Note that then each subset of $j\leq n+1$ vertices of a generic spherical polyhedron is affinely independent and, if $n\geq 3$, $\mathcal{K}$ does not have two parallel edges, because otherwise $\mathcal{K}$ would have $4\leq n+1$ coplanar points and they would be affinely dependent.

Lemma 4.7. *Let $\mathcal{K}'$ be the spherical polyhedron whose set of vertices is $\mathfrak{F}':=\{p'_1,\dots,p'_k\}$ and let $0<\varepsilon<1$. Then there is a generic spherical polyhedron $\mathcal{K}$ whose set of vertices is $\mathfrak{F}:=\{p_1,\dots,p_k\}$ such that $p_i\in\mathcal{B}_n(p'_i,\varepsilon)$ for each $i=1,\dots,k$.*

Proof. Consider the pure dimensional irreducible real algebraic variety

$$\mathbb{T}:=\mathbb{S}^{n-1}\times\overset{(k)}{\cdots}\times\mathbb{S}^{n-1}\subset\mathbb{R}^n\times\overset{(k)}{\cdots}\times\mathbb{R}^n\equiv\mathbb{R}^{kn}.$$

For each spherical polyhedron $\mathcal{K}\subset\mathbb{R}^n$ with vertices $\{q_1,\dots,q_k\}$, we associate (uniquely up to the order of the vertices) the point $q:=(q_1,\dots,q_k)\in\mathbb{T}$; obviously, not all the points of $\mathbb{T}$ define spherical polyhedra because we require from them to be n-dimensional and to have distinct vertices. Define on $\mathbb{T}$ the metric

$$d(q,q')=\sqrt{\textstyle\sum_{i=1}^k d^2(q_i,q'_i)},\quad q:=(q_1,\dots,q_k),q':=(q'_1,\dots,q'_k)\in\mathbb{T}.$$

In particular, if $d(q,q')<\varepsilon$, then $d(q_i,q'_i)<\varepsilon$ for $i=1,\dots,k$. Let us show that the set $\mathbb{G}$ of all points of $\mathbb{T}$ corresponding to generic spheric polyhedra is dense in $\mathbb{T}$. Since $\mathbb{T}$ is irreducible and pure dimensional, it is enough to check that $Z:=\mathbb{T}\setminus\mathbb{G}$ is a proper algebraic subset of $\mathbb{G}$. Note that Z is the set of points $q:=(q_1,\dots,q_k)\in\mathbb{T}$ such that there is a subset of $n+1$ points affinely dependent. Write $\mathtt{x}:=(\mathtt{x}_1,\dots,\mathtt{x}_k)$, $\mathtt{x}_i:=(\mathtt{x}_{i1},\dots,\mathtt{x}_{in})$ and for each set $I:=\{i_1,\dots,i_{n+1}\}\subset\{1,\dots,k\}$ consider the polynomial

$$F_I(\mathtt{x}):=\det\begin{pmatrix}1&\mathtt{x}_{i_11}&\cdots&\mathtt{x}_{i_1n}\\ \vdots&\vdots&\ddots&\vdots\\ 1&\mathtt{x}_{i_j1}&\cdots&\mathtt{x}_{i_jn}\end{pmatrix}$$

Clearly, the points $q_{i_1},\dots,q_{i_n}$ are affinely dependent if and only if $F_I(q)=0$ and so $Z=\{\prod_I F_I(x)=0\}\cap\mathbb{T}$ is an algebraic set. Moreover, such subset is proper because $\mathbb{G}$ is nonempty. In fact, to construct a point q of $\mathbb{G}$, we can start by choosing $n+1$ affinely independent points of $\mathbb{S}^{n-1}$, adding successively new points of $\mathbb{S}^{n-1}$ (until we reach k) in such a way that, once we have chosen $\{q_1,\dots,q_r\}$ with $r\geq n+1$, we pick $q_{r+1}\in\mathbb{S}^{n-1}$ not belonging to any of the hyperplanes generated by subsets of n points of $\{q_1,\dots,q_r\}$.

Finally, the ball $\mathcal{B}_{nk}(p',\varepsilon)$ contains points of $\mathbb{G}$, and so $p:=(p_1,\dots,p_k)\in\mathbb{G}\cap\mathcal{B}_{nk}(p',\varepsilon)$ provides a generic spherical polyhedron $\mathcal{K}$ whose vertices $p_i\in\mathcal{B}_n(p'_i,\varepsilon)$. □

Lemma 4.8. *For each $0<r<1$ there is a generic spherical polyhedron $\mathcal{K}\subset\mathbb{R}^n$ such that $\mathcal{B}_n(\mathbf{0},r)\subset\mathcal{K}$.*

Proof. Denote $\varepsilon':=\frac{1}{2}\sqrt{2(1-r)}$ and consider the open covering $\{\mathcal{B}_n(p',\varepsilon'):\ p'\in\mathbb{S}^{n-1}\}$ of the sphere $\mathbb{S}^{n-1}$. Since $\mathbb{S}^{n-1}$ is compact, we can extract a finite subcovering $\{\mathcal{B}_n(p'_i,\varepsilon'):1\leq i\leq k\}$. By Lemma 4.7 there are points $p_1,\dots,p_k\in\mathbb{S}^{n-1}$ such that $d(p_i,p'_i)<\varepsilon'$ for $i=1,\dots,k$ and whose convex hull is a generic spherical polyhedron.

Next, write $\varepsilon:=2\varepsilon'$ and observe that $\{\mathcal{B}_n(p_i,\varepsilon):i=1,\dots,k\}$ is an open covering of $\mathbb{S}^{n-1}$, because $\mathcal{B}_n(p'_i,\varepsilon')\subset\mathcal{B}_n(p_i,\varepsilon)$. We claim that if $\mathcal{F}$ is an arbitrary facet of the polyhedron $\mathcal{K}$ and H is the hyperplane generated by $\mathcal{F}$, then

$$d(\mathbf{0},H)\geq r=1-\frac{\varepsilon^2}{2}.$$

Indeed, suppose by way of contradiction $d(\mathbf{0},H)=d<r$. We may assume $H:=\{x_1=d\}$ and $\mathcal{K}\subset H^+:=\{x_1\leq d\}$. Let $q:=(1,0,\dots,0)\in\mathbb{S}^{n-1}$ and let $p_i=(p_{i1},\dots,p_{in})\in\mathbb{S}^{n-1}$ be a vertex of $\mathcal{K}$; it holds that $\sum_{j=1}^n p_{ij}^2=1$ and $p_{i1}\leq d$. Thus,

$$d(p_i,q)=\sqrt{(p_{i1}-1)^2+p_{i2}^2+\cdots+p_{in}^2}=\sqrt{2(1-p_{i1})}>\sqrt{2(1-r)}=\varepsilon.$$

But then $q\notin\mathcal{B}_n(p_i,\varepsilon)$ for $i=1,\dots,k$, a contradiction. We conclude that $d(\mathbf{0},H)\geq r$.

Now, since $\mathcal{K}=\bigcap_{i=1}^r H_i^+$, where each H_i^+ is the half-space of $\mathbb{R}^n$ that contains $\mathcal{K}$ and whose boundary H_i contains a facet of $\mathcal{K}$, and moreover $\mathcal{B}_n(\mathbf{0},r)\subset H_i^+$ for each $i=1,\dots,r$, we deduce that $\mathcal{B}_n(\mathbf{0},r)\subset\mathcal{K}$. □

Lemma 4.9. *Let $m>n$ be a positive integer. Then there is $r>0$ such that if $\mathcal{K}\subset\mathbb{R}^n$ is a convex polyhedron with $\overline{\mathcal{B}}_n(\mathbf{0},r)\subset\mathcal{K}\subset\overline{\mathcal{B}}_n(\mathbf{0},1)$, it holds that $\mathcal{K}$ has more than m vertices.*

Proof. First, let $\varepsilon>0$ and $r:=1-\varepsilon$ and pick two points $p_1,p_2\in\overline{\mathcal{B}}_n(\mathbf{0},1)\setminus\overline{\mathcal{B}}_n(\mathbf{0},1-\varepsilon)$ such that the line ℓ passing through p_1 and p_2 does not intersect $\overline{\mathcal{B}}_n(\mathbf{0},r)$. We claim that $d(p_1,p_2)\leq 2\sqrt{2\varepsilon-\varepsilon^2}$.

Indeed, let $q_0\in\ell$ be the closest point of ℓ to the origin and let q_1,q_2 be the intersection points of ℓ with $\mathbb{S}^{n-1}$. Clearly, $d(p_1,p_2)\leq d(q_1,q_2)$ and the line passing through $\mathbf{0}$ and q_0 is perpendicular to ℓ. In fact,

$$1=d(\mathbf{0},q_1)^2=d(\mathbf{0},q_0)^2+d(q_0,q_1)^2\geq(1-\varepsilon)^2+d(q_0,q_1)^2\implies d(q_0,q_1)\leq\sqrt{2\varepsilon-\varepsilon^2}.$$

Thus, $d(p_1,p_2)\leq d(q_1,q_2)=2d(q_0,q_1)=2\sqrt{2\varepsilon-\varepsilon^2}$, as claimed.

Let now $0<\varepsilon<1/4$ be such that $2(m-1)\sqrt{2\varepsilon-\varepsilon^2}<1/4$ and $r:=1-\varepsilon$. Suppose, by way of contradiction that there is a convex polyhedron $\mathcal{K}\subset\mathbb{R}^n$ with at most m vertices such that $\overline{\mathcal{B}}_n(\mathbf{0},r)\subset\mathcal{K}\subset\overline{\mathcal{B}}_n(\mathbf{0},1)$. After an orthogonal change of coordinates, we may assume that one of the vertices of $\mathcal{K}$ is $(0,\dots,0,t)$ for some $r<t\leq 1$. Let $\mathcal{A}$ be an edge of $\mathcal{K}$ and let ℓ be the line generated by $\mathcal{A}$. Since $\mathcal{A}\subset\partial\mathcal{K}$, we have that $\ell\cap\overline{\mathcal{B}}_n(\mathbf{0},r)=\varnothing$ and as we have proved above the length of $\mathcal{A}$ is smaller than $2\sqrt{2\varepsilon-\varepsilon^2}$. Thus, since $\mathcal{K}$ has at most m vertices, the distance between two vertices of $\mathcal{K}$ is at most $2(m-1)\sqrt{2\varepsilon-\varepsilon^2}<1/4$; hence, since one of the vertices of $\mathcal{K}$ is $(0,\dots,0,t)$ and $t>3/4$, we deduce that all the vertices of $\mathcal{K}$ (and so $\mathcal{K}$) are contained in the half-ball $\overline{\mathcal{B}}_n(\mathbf{0},1)\cap\{x_n\geq 0\}$, which contradicts the fact that $\overline{\mathcal{B}}_n(\mathbf{0},r)\subset\mathcal{K}$. □

4.3. **Proof of Theorem 4.2.** Let r be the positive integer of Lemma 4.9 for $m:=n+1$ and let $\mathcal{K}\subset\mathbb{R}^n$ be a generic spherical polyhedron such that $\mathcal{B}_n(\mathbf{0},r)\subset\mathcal{K}$ (see Lemma 4.8). Consider the projection $\pi:\mathbb{R}^n\to\mathbb{R}^n$ in the direction of the line $\vec{\ell}$ onto the hyperplane H and denote $\mathcal{P}:=\pi(\mathcal{K})$. Let H' be the hyperplane perpendicular to ℓ that contains the origin. After an orthogonal change of coordinates, we may assume that $\vec{\ell}$ is generated by the vector $\vec{e}_n:=(0,\dots,0,1)$, and $H':=\{x_n=0\}$. Let

$$\pi':\mathbb{R}^n\to H',\ x:=(x_1,\dots,x_{n-1},x_n)\to(x',0):=(x_1,\dots,x_{n-1},0)$$

be the projection onto H' in the direction of $\vec{\ell}$ and denote

$$\mathcal{P}':=\pi(\mathcal{K}),\quad \mathcal{P}'_0:=\{x'\in\mathbb{R}^{n-1}:\ (x',0)\in\mathcal{P}'\}.$$

Notice that ${\pi'}^{-1}(\mathcal{P}') := \mathcal{P}'_0 \times \mathbb{R}$ and $\mathcal{P} := \pi(\mathcal{K}) = (\mathcal{P}'_0 \times \mathbb{R}) \cap H$. Thus, $\mathcal{P}$ can be considered as the graph of an affine function over $\mathcal{P}'_0$; hence, it is affinely equivalent to $\mathcal{P}'_0$. Moreover, we have $\pi(\overline{\mathcal{B}}_n(\mathbf{0}, s)) = \overline{\mathcal{B}}_{n-1}(\mathbf{0}, s) \times \{0\}$ for each $s > 0$.

Suppose now that $\mathcal{K} \cap H = \mathcal{P}$. Since $\mathcal{K}$ is a generic spherical polyhedron, each hyperplane of $\mathbb{R}^n$ contains at most n vertices of $\mathcal{K}$ and $\mathcal{K}$ has nonparallel edges. Thus, by Lemma 4.4 (ii), the polyhedron $\mathcal{P} = \mathcal{K} \cap H$ has at most $n + 1$ vertices.

On the other hand, the convex polyhedron $\mathcal{P}'_0$ satisfies $\overline{\mathcal{B}}_{n-1}(\mathbf{0}, r) \subset \mathcal{P}' \subset \overline{\mathcal{B}}_{n-1}(\mathbf{0}, 1)$ and, by Lemma 4.9, it has at least $n + 2$ vertices. But this is against the fact that $\mathcal{P}$ and $\mathcal{P}'_0$ are affinely equivalent; we conclude that $\mathcal{K}$ has no sectional projections. □

REFERENCES

[Be1] M. Berger: Geometry. I. *Universitext.* Springer-Verlag, Berlin: 1987.

[Be2] M. Berger: Geometry. II. *Universitext.* Springer-Verlag, Berlin: 1987.

[BCR] J. Bochnak, M. Coste, M.F. Roy: Real algebraic geometry. *Ergeb. Math.* **36**. Springer-Verlag, Berlin: 1998.

[Fe] J.F. Fernando: On the one dimensional polynomial and regular images of $\mathbb{R}^n$. *Preprint* RAAG (2012). `http://www.mat.ucm.es/~josefer/pdfs/preprint/curvepolregim.pdf`

[FG1] J.F. Fernando, J.M. Gamboa: Polynomial images of $\mathbb{R}^n$. *J. Pure Appl. Algebra* **179** (2003), no. 3, 241–254.

[FG2] J.F. Fernando, J.M. Gamboa: Polynomial and regular images of $\mathbb{R}^n$. *Israel J. Math.* **153** (2006), 61–92.

[FGU] J.F. Fernando, J.M. Gamboa, C. Ueno: On convex polyhedra as regular images of $\mathbb{R}^n$. *Proc. London Math. Soc.* (3) **103** (2011), 847-878.

[FU1] J.F. Fernando, C. Ueno: On the set of infinite points of a polynomial image of $\mathbb{R}^n$. *Preprint* RAAG (2011). `http://www.mat.ucm.es/~josefer/pdfs/preprint/comppol-dimn.pdf`

[FU2] J.F. Fernando, C. Ueno: On complements of convex polyhedra as polynomial and regular images of $\mathbb{R}^n$. *Preprint* RAAG (2012).

[R] T.R. Rockafellar: Convex analysis. *Princeton Mathematical Series*, **28**. Princeton University Press, Princeton, N.J: 1970.

[U1] C. Ueno: A note on boundaries of open polynomial images of $\mathbb{R}^2$. *Rev. Mat. Iberoam.* **24** (2008), no. 3, 981-988.

[U2] C. Ueno: On convex polygons and their complements as images of regular and polynomial maps of $\mathbb{R}^2$. *J. Pure Appl. Algebra* **216**, no. 11, 2436-2448.

[U3] C. Ueno: Convex Polygons via Polynomial Origami. *Preprint* RAAG (2012).

[Z] G.M. Ziegler: Lectures on Polytopes. *Graduate Texts in Mathematics* **152**. Springer-Verlag, New-York: 1995

Departamento of Álgebra, Facultad of Ciencias Matemáticas, Universidad Complutense of Madrid, 28040 MADRID (SPAIN)

E-mail address: `josefer@mat.ucm.es`

Departamento de Matemáticas, IES La Vega de San José, Paseo de San José, s/n, Las Palmas de Gran Canaria, 35015 LAS PALMAS (SPAIN).

E-mail address: `carlos.ueno@terra.es`